

13 FEB 2001

## CERTIFICATION UNDER 37 CFR 1.10

E1 685061392US

Express Mail Mailing Number

February 13, 2001

Date of Deposit

I hereby certify that this paper or fee, and any documents referred to as enclosed herein, are being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents, Box PCT, Washington, D.C. 20231.

Cedric Rodgers

Name of Person Mailing Application

Signature of Person Mailing Application

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

VOS-101

TRANSMITTAL LETTER TO THE UNITED STATES  
DESIGNATED/ELECTED OFFICE (DO/EO/US)  
CONCERNING A FILING UNDER 35 U.S.C. 371

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)  
Not known: 09/762767

INTERNATIONAL APPLICATION NO.  
PCT/DE99/02601

INTERNATIONAL FILING DATE  
16 August 1999

PRIORITY DATE CLAIMED  
14 August 1998

## TITLE OF INVENTION

ISOLATED AND PURIFIED HUMAN SOLUBLE GUANYLYL CYCLASE  $\alpha 1/\beta 1$  (hSGCa1/ $\beta 1$ )

## APPLICANT(S) FOR DO/EO/US

Harald SCHMIDT, Ulrike ZABEL and Wolfgang POLLER

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☐ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☒ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
  - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
  - b. ☒ has been transmitted by the International Bureau.
  - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US)
6. ☒ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
  - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
  - b. ☐ have been transmitted by the International Bureau.
  - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
  - d. ☒ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An oath or declaration of the inventor(s) 35 U.S.C. 371(c)(4)) and **Power of Attorney** (Unexecuted).
10. ☒ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

## Items 11 to 16 below concern other document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☐ A FIRST preliminary amendment.
- ☐ A SECOND or SUBSEQUENT preliminary amendment.

US APPLICATION NO. (if known, see 37 CFR 1.5) <div style="font-size: 1.5em; font-weight: bold; margin-top: 5px;">097762767</div>	INTERNATIONAL APPLICATION NO. PCT/DE99/02601	ATTORNEYS DOCKET NUMBER VOS-101																																								
14. <input type="checkbox"/> A substitute specification and claims amended to conform the translated application to US practice 15. <input type="checkbox"/> A change of power of attorney and/or address letter. 16. <input checked="" type="checkbox"/> Other items or information: (a) copy of the First Page of International Publication; (b) copy of International Search Report; (c) copies of Written Opinion and International Preliminary Examination Report																																										
17. <input checked="" type="checkbox"/> The following fees are submitted:  <b>BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5)):</b>  Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO ..... \$1,000.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO ..... \$860.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search fee (37 CFR 1.445(a)(2)) paid to USPTO .... \$710.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4) ..... \$690.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4) ..... \$100.00  <div style="text-align: right; font-weight: bold;">ENTER APPROPRIATE BASIC FEE AMOUNT =</div>	<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <th style="text-align: left; padding: 2px;">CALCULATIONS</th> <th style="text-align: left; padding: 2px;">PTO USE ONLY</th> </tr> <tr> <td style="height: 100px; vertical-align: top; padding: 2px;"> <div style="text-align: right; font-weight: bold;">\$860.00</div> </td> <td></td> </tr> <tr> <td style="padding: 2px;">           Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).         </td> <td style="text-align: right; padding: 2px;">\$0.00</td> </tr> <tr> <td style="padding: 2px;"> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 20%;">Claims</th> <th style="width: 20%;">Number Filed</th> <th style="width: 20%;">Number Extra</th> <th style="width: 40%;">Rate</th> </tr> <tr> <td>Total Claims</td> <td style="text-align: center;">19</td> <td style="text-align: center;">0</td> <td style="text-align: right;">\$18.00</td> </tr> <tr> <td>Independent Claims</td> <td style="text-align: center;">6</td> <td style="text-align: center;">0</td> <td style="text-align: right;">\$80.00</td> </tr> </table> </td> <td style="text-align: right; padding: 2px;">\$240.00</td> </tr> <tr> <td style="padding: 2px;">           MULTIPLE DEPENDENT CLAIM(S) (if applicable)         </td> <td style="text-align: right; padding: 2px;">\$270.00</td> </tr> <tr> <td style="padding: 2px;"> <div style="text-align: right; font-weight: bold;">TOTAL OF ABOVE CALCULATIONS =</div> </td> <td style="text-align: right; padding: 2px;">\$1,370.00</td> </tr> <tr> <td style="padding: 2px;"> <input type="checkbox"/> Small Entity Status: Applicant claims small entity status. (See 37 CFR 1.27.) Reduction of 1/2 for filing by small entity (if applicable).         </td> <td style="text-align: right; padding: 2px;">\$0.00</td> </tr> <tr> <td style="padding: 2px;"> <div style="text-align: right; font-weight: bold;">SUBTOTAL =</div> </td> <td style="text-align: right; padding: 2px;">\$1,370.00</td> </tr> <tr> <td style="padding: 2px;">           Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).         </td> <td style="text-align: right; padding: 2px;">\$0.00</td> </tr> <tr> <td style="padding: 2px;"> <div style="text-align: right; font-weight: bold;">TOTAL NATIONAL FEE =</div> </td> <td style="text-align: right; padding: 2px;">\$1,370.00</td> </tr> <tr> <td style="padding: 2px;">           Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) \$40 per property         </td> <td style="text-align: right; padding: 2px;">\$0.00</td> </tr> <tr> <td style="padding: 2px;"> <div style="text-align: right; font-weight: bold;">TOTAL FEES ENCLOSED =</div> </td> <td style="text-align: right; padding: 2px;">\$1,370.00</td> </tr> <tr> <td style="padding: 2px;"></td> <td style="text-align: right; padding: 2px;">Amount to be refunded \$</td> </tr> <tr> <td style="padding: 2px;"></td> <td style="text-align: right; padding: 2px;">charged \$</td> </tr> </table>		CALCULATIONS	PTO USE ONLY	<div style="text-align: right; font-weight: bold;">\$860.00</div>		Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).	\$0.00	<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 20%;">Claims</th> <th style="width: 20%;">Number Filed</th> <th style="width: 20%;">Number Extra</th> <th style="width: 40%;">Rate</th> </tr> <tr> <td>Total Claims</td> <td style="text-align: center;">19</td> <td style="text-align: center;">0</td> <td style="text-align: right;">\$18.00</td> </tr> <tr> <td>Independent Claims</td> <td style="text-align: center;">6</td> <td style="text-align: center;">0</td> <td style="text-align: right;">\$80.00</td> </tr> </table>	Claims	Number Filed	Number Extra	Rate	Total Claims	19	0	\$18.00	Independent Claims	6	0	\$80.00	\$240.00	MULTIPLE DEPENDENT CLAIM(S) (if applicable)	\$270.00	<div style="text-align: right; font-weight: bold;">TOTAL OF ABOVE CALCULATIONS =</div>	\$1,370.00	<input type="checkbox"/> Small Entity Status: Applicant claims small entity status. (See 37 CFR 1.27.) Reduction of 1/2 for filing by small entity (if applicable).	\$0.00	<div style="text-align: right; font-weight: bold;">SUBTOTAL =</div>	\$1,370.00	Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).	\$0.00	<div style="text-align: right; font-weight: bold;">TOTAL NATIONAL FEE =</div>	\$1,370.00	Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) \$40 per property	\$0.00	<div style="text-align: right; font-weight: bold;">TOTAL FEES ENCLOSED =</div>	\$1,370.00		Amount to be refunded \$		charged \$
CALCULATIONS	PTO USE ONLY																																									
<div style="text-align: right; font-weight: bold;">\$860.00</div>																																										
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).	\$0.00																																									
<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 20%;">Claims</th> <th style="width: 20%;">Number Filed</th> <th style="width: 20%;">Number Extra</th> <th style="width: 40%;">Rate</th> </tr> <tr> <td>Total Claims</td> <td style="text-align: center;">19</td> <td style="text-align: center;">0</td> <td style="text-align: right;">\$18.00</td> </tr> <tr> <td>Independent Claims</td> <td style="text-align: center;">6</td> <td style="text-align: center;">0</td> <td style="text-align: right;">\$80.00</td> </tr> </table>	Claims	Number Filed	Number Extra	Rate	Total Claims	19	0	\$18.00	Independent Claims	6	0	\$80.00	\$240.00																													
Claims	Number Filed	Number Extra	Rate																																							
Total Claims	19	0	\$18.00																																							
Independent Claims	6	0	\$80.00																																							
MULTIPLE DEPENDENT CLAIM(S) (if applicable)	\$270.00																																									
<div style="text-align: right; font-weight: bold;">TOTAL OF ABOVE CALCULATIONS =</div>	\$1,370.00																																									
<input type="checkbox"/> Small Entity Status: Applicant claims small entity status. (See 37 CFR 1.27.) Reduction of 1/2 for filing by small entity (if applicable).	\$0.00																																									
<div style="text-align: right; font-weight: bold;">SUBTOTAL =</div>	\$1,370.00																																									
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).	\$0.00																																									
<div style="text-align: right; font-weight: bold;">TOTAL NATIONAL FEE =</div>	\$1,370.00																																									
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) \$40 per property	\$0.00																																									
<div style="text-align: right; font-weight: bold;">TOTAL FEES ENCLOSED =</div>	\$1,370.00																																									
	Amount to be refunded \$																																									
	charged \$																																									

US APPLICATION NO. (if known, see 37 CFR 1.51)

091762767

INTERNATIONAL APPLICATION NO.  
PCT/DE99/02601

ATTORNEYS DOCKET NUMBER  
VOS-101

- a. ☒ A check in the amount of \$1,370.00 to cover the above fees is enclosed.
- b. ☐ Please charge my Deposit Account No. \_\_\_\_\_ in the amount of \_\_\_\_\_ to cover the above fees. A duplicate copy of this sheet is enclosed.
- c. ☒ The Commissioner is hereby authorized to charge any additional filing fees which may be required, or credit any overpayment to Deposit Account No. 15-0508. A duplicate copy of this sheet is enclosed.

**NOTE:** Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

OLSON & HIERL, LTD.  
20 North Wacker Drive  
36th Floor  
Chicago, Illinois 60606

Telephone: (312) 580-1180  
Customer No.: 002387



SIGNATURE

Talivaldis Cepuritis

NAME

20,818

REGISTRATION NO.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE  
 UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US)

Applicant(s): Harald Schmidt, et al. )  
 )  
 Serial No.: 09/762,767 )  
 )  
 Filed: February 13, 2001 )  
 )  
 For: ISOLATED AND PURIFIED HUMAN )  
 SOLUBLE GUANYLYL CYCLASE )  
 $\alpha 1 / \beta 1$  (HsGC  $\alpha 1 / \beta 1$ ) ) Atty. Docket No.: VOS-101

5000

COMMUNICATION

Box PCT  
 Commissioner for Patents  
 Washington, D.C. 20231

Sir:


In response to the Notification of a Defective Response dated 20 September 2001, we submit the following in compliance with 37 CFR 1.821-1.825:

- (a) New paper copy of Sequence Listing; and
- (b) Computer-readable form of the Sequence Listing.

These submissions do not constitute new matter and are supported in the application as filed. To the best of my information and belief, the sequence listing information recorded in computer-readable form is identical to the paper copy of the sequence listing.

Please charge any additional fees concerning this matter or credit any overpayment to our Deposit Account No. 15-0508.

Respectfully submitted,

By   
 Talivaldis Cepuritis (Reg. No. 20,818)

OLSON & HIERL, LTD.  
 20 North Wacker Drive  
 36th Floor  
 Chicago, Illinois 60606  
 (312) 580-1180

Serial No. 09/762,767 - - - - - 2


CERTIFICATION UNDER 37 CFR 1.10

"Express Mail" mailing label number: **EL 841695088 US**

I hereby certify that this correspondence, together with any other documents and/or fees referred to as enclosed herein, is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service on October 17, 2001 and is addressed to: Box PCT, Commissioner for Patents, Washington, D.C. 20231.

Cedric Rodgers

(Typed or printed name of person mailing paper or fee)



(Signature of person mailing paper or fee)

**Isolated and purified human soluble guanylyl cyclase  $\alpha 1/\beta 1$   
(hsGC  $\alpha 1/\beta 1$ )**

**The technical field of the invention**

5

The present invention relates to the expression of the cDNA clones for the subunits  $\alpha 1$  (hsGC $\alpha 1$ ) and  $\beta 1$  (hsGC $\beta 1$ ) of human soluble guanylyl cyclase and the subsequent purification of the active enzyme and the use thereof, the medical application of the expression of these clones by gene transfer, as well as antibodies to peptides derived from the sequence and the use thereof.

10

**State of the art**

- 15 The endogenous NO/cGMP signaling system mediates important functions such as vasodilation, inhibition of platelet aggregation, neurotransmission, and the immune response. In addition, it is involved in the development of various disease states such as ischemia-reperfusion and inflammatory injuries (Schmidt and Walter, 1994). Therefore, the NO/cGMP system has long been an important starting point for the
- 20 development of novel drugs for the therapy of coronary heart disease, susceptibility to thrombosis, cardiac insufficiency, angina pectoris, cardiac-dependent pulmonary oedema, hypertensive crises, inflammatory states and cardiac infarction. Until now, such therapies have employed various so-called NO donors, e.g. nitroglycerin, that release NO, thereby replacing endogenous NO and activating soluble guanylyl
- 25 cyclase (sGC) (Fig. 1). sGC forms cGMP, which mediates the effects of the NO/cGMP pathway via various intracellular receptor enzymes. The application of NO donors has two limitations: I) The repeated application of NO donors results in tolerance in the patient, i.e. loss of activity, when reappplied; II) NO reacts with  $O_2^-$  to form peroxynitrite, which is cytotoxic and less effective in activating sGC. Thus, direct
- 30 activation of sGC by novel, non-NO-containing pharmaceuticals for therapy or gene transfer of sGC would be desirable strategies. Non-NO-releasing activators of sGC are potentially lacking tolerance and NO toxicitation. Recently, "YC-1" (3-(5'-hydroxymethyl-2'-furyl)-1-benzylindazol) has been described as the first non-NO-releasing activator of sGC in platelets (Ko et al., 1994; Wu et al., 1995). YC-1 also

activates purified sGC from bovine lung and potentiates the activation of NO (Friebe et al., 1996). The effect on human sGC has not been investigated. The human isoform of sGC is not yet available for pharmacological screening.

- 5 The NO receptor sGC consists of two subunits,  $\alpha$  and  $\beta$ , which together form an enzymatically active heterodimer. Three different isoforms of the  $\alpha$  and three of the  $\beta$  subunits have been described in the literature, although from different species. The best investigated isoforms, the bovine  $\alpha 1/\beta 1$  and the rat  $\alpha 1/\beta 1$  isoforms, have particular significance for cardiovascular research. Until recently, a human
- 10 homologue of the bovine and rat sGC (e.g. sGC $\alpha 1$ /sGC $\beta 1$  heterodimers) was unknown. cDNA sequences have been published that reportedly correspond to a human sGC $\alpha 3$  and  $\beta 3$  isoform (Giulli et al., 1992). While sGC $\beta 3$  showed a high homology to sGC $\beta 1$  (bovine/rat), the sGC $\alpha 3$  sequence contained two restricted regions without homology to sGC $\alpha 1$  (bovine/rat), here designated S1 and S2 (Fig. 2).
- 15 In addition, it was unknown whether sGC $\alpha 3$  and sGC $\beta 3$  can form a functional sGC heterodimer and which role the S1 and S2 regions play. Furthermore, no other human sGC subunit has yet been expressed as a protein. Recently, a sequence designated hsGC $\alpha 1$  was published in the GeneBank (Accession No. U58855) that lacks the sequence differences to sGC $\alpha 1$  from bovine and rat tissues in the regions
- 20 S1 and S2. In addition, an alternatively spliced product of hsGC $\beta 3$  was recently published in the GeneBank (Accession No. AF020340) that was designated by the authors as an alternatively spliced form of hsGC $\beta 1$ . The physiological significance of this splice variant of hsGC $\beta 1/3$  is unclear. Thus, the question arises as to which of these isoforms is responsible for which physiological function in which cell types.

25

- Furthermore, no antibodies to the human sGC $\alpha 1/\beta 1$  are currently available that are monospecific, directed against the human sequences, or that have been shown to be suitable for immunoblots with human tissues. Peptide antibodies reported thus far only partially show these features: Harteneck et al. and Guthmann et al. used a
- 30 peptide sequence (VYKVETVGD KYMTVSGLP) that is highly conserved in guanylyl cyclases. Therefore, cross-reaction with particulate guanylyl cyclases (e.g. GC-C) can be expected. Guthmann et al. used a peptide sequence (YGPEVVEDIKKEA) identical to hsGC $\beta 1$  and a peptide sequence (KKDVEEANANFLGKASGID) identical

to hSGC $\alpha$ 1 except for two amino acid exchanges. However, the function of these antibodies in immunoblots was only shown for *enriched* hSGC from human platelets. In addition, the antisera to hSGC $\alpha$ 1 detect a second, unspecific product. Humbert et al. and Koesling et al. used a peptide sequence (SRKNTGTEETE QDEN) of bovine sGC $\beta$ 1 that partially (amino acids 1-10) overlaps with the peptide used here for the human sGC $\beta$ 1 (amino acids 13-22) and is identical in this region. The C-terminus of this bovine peptide (amino acids 11-15), however, is clearly different from the human sequence. On the other hand, the antiserum to this peptide was not tested with a human protein but rather only used for immunoprecipitation of the bovine sGC.

Thus, for the human isoform  $\alpha$ 1/ $\beta$ 1 which is important for cardiovascular research, neither *native* protein nor *recombinant* protein, nor a one-step purification protocol, nor specific antibodies have been available. An approach for gene therapy (e.g. with adenoviruses or the like) has also not been described yet.

### The technical problem of the invention

The NO-independent activation of the human sGC $\alpha$ 1/ $\beta$ 1 is a promising approach to finding novel drugs or gene transfer techniques for cardiovascular therapy that neither result in tolerance in the patient nor form cytotoxic peroxynitrite. To find such pharmaceuticals, a mass screening for suitable active substances is necessary. Such a pharmacological screening for specific activators or inhibitors via animal testing is too expensive and is not reasonable due to species differences, possible side effects, and effects on other isoforms. Cell culture systems have the disadvantage that only with substantial additional effort can it be determined at which points of the signaling cascade substances are effective. In addition, cell culture is expensive and touchy. The purification of a protein from animal tissues is labor-intensive and results in a low yield. More importantly, because of species differences the results of such a screening are not generally transferable. Here, in particular, the question arises of the significance of the isoforms hSGC $\alpha$ 1 and hSGC $\alpha$ 3 and whether hSGC $\beta$ 3 in fact corresponds to a human hSGC $\beta$ 1. This is an important factor to consider in selecting the appropriate target protein for pharmacological screening or gene therapy. For optimal screening, the human homologue of the protein sGC $\alpha$ 1/ $\beta$ 1 would have to be



purified in large amounts and to be available at low cost. A purification of large amounts of human native sGC $\alpha$ 1/ $\beta$ 1 protein from human tissues is not possible. Thus, other methods are needed for the drug screening protocol. In addition, no purified antibodies are available for the detection of human sGC $\alpha$ 1/ $\beta$ 1 that have been shown to be suitable for diagnostic uses, e.g. in normal immunoblotting, ELISA, RIA, or EIA, among other techniques. An approach for the artificial expression of hsGC by gene transfer in humans that could be used therapeutically is also not available.

Therefore, the present invention is based on the technical problem of providing isolated and purified human sGC $\alpha$ 1/ $\beta$ 1 as well as a process for its production and purification. In addition, a further technical problem of the present invention is to provide antibodies directed against human sGC $\alpha$ 1/ $\beta$ 1. A further technical problem is the provision of expression vectors containing the cDNA of human sGC $\alpha$ 1/ $\beta$ 1 based on adenoviruses. Finally, the technical problem of the present invention is the provision of human sGC $\alpha$ 1/ $\beta$ 1 in purified form and in manageable amounts for drug screening assays aimed at detecting modulators, inhibitors, and activators of human sGC $\alpha$ 1/ $\beta$ 1.

#### **The solution of the technical problem**

The solution to the above technical problem is provided by the subject-matter of the claims and the following description of the invention.

An object of the invention is a human soluble guanylyl cyclase  $\alpha$ 1/ $\beta$ 1 (hsGC $\alpha$ 1/ $\beta$ 1) in an isolated form purified to apparent homogeneity.

Another object of the invention is a process for the production of subunit  $\alpha$ 1 and/or  $\beta$ 1 of human soluble guanylyl cyclase comprising the expression of expression vectors containing the DNA sequence for hsGC $\alpha$ 1 and/or hsGC $\beta$ 1 in prokaryotic and eukaryotic host cells, and obtaining of the subunit or subunits.

In a preferred embodiment of the process of the present invention for the production of subunit  $\alpha$ 1 and/or  $\beta$ 1 of human soluble guanylyl cyclase, the step of obtaining the

subunit or subunits comprises the lysis of cells, affinity chromatography of the cell lysate, and finally, the elution of the subunit or subunits.

- 5 In a further preferred embodiment of the process of the present invention for the production of subunit  $\alpha 1$  and/or  $\beta 1$  of human soluble guanylyl cyclase, the expression vector contains additionally at least one DNA sequence of a domain for specific affinity chromatography (affinity tag) with attached protease cleavage site.

- 10 Another object of the present invention is a process for the production of human soluble guanylyl cyclase  $\alpha 1/\beta 1$  (hsGC $\alpha 1/\beta 1$ ) comprising the separate expression of an expression vector containing the DNA sequence for hsGC $\alpha 1$  or hsGC $\beta 1$  in prokaryotic or eukaryotic host cells, obtaining the subunits, and combining the subunits hsGC $\alpha 1$  and hsGC $\beta 1$  into the dimeric guanylyl cyclase  $\alpha 1/\beta 1$  (hsGC $\alpha 1/\beta 1$ ).

- 15 A further preferred embodiment of the process of the present invention for the production of human soluble guanylyl cyclase  $\alpha 1/\beta 1$  (hsGC $\alpha 1/\beta 1$ ) also comprises the coexpression of the DNA sequences for hsGC $\alpha 1$  and hsGC $\beta 1$  in prokaryotic or eukaryotic host cells, the lysis of the cells containing hsGC $\alpha 1$  and hsGC $\beta 1$ , affinity chromatography, and the subsequent elution of hsGC $\alpha 1/\beta 1$ .

20

- Another object of the present invention is the use of a nucleic acid sequence that codes for subunits hsGC $\alpha 1$  and/or hsGC $\beta 1$  of human soluble guanylyl cyclase  $\alpha 1/\beta 1$  for somatic gene therapy, particularly for the prevention and therapy of atherosclerosis and its complications, restenosis, ischemia (infarction), peripheral occlusive diseases, and arterial hypertension, as well as, in patients with risk factors, for the prevention of atherosclerosis, transient ischemic attacks, cerebral ischemia, stroke, coronary heart disease, status post coronary bypass grafting, carotid stenosis, heart insufficiency and liver dysfunction, and as a supplement to therapy with sGC activators, sGC-sensitizing substances, NO donors, or phosphodiesterase inhibitors.
- 25
- 30

In a particularly preferred embodiment, adenoviral vectors containing hsGC $\alpha 1$  or hsGC $\beta 1$  cDNA are used in somatic gene therapy. Other vector systems, however, can also be applied for gene therapeutic, medicinal use.

In a preferred embodiment, adenoviral Ad5 and other suitable vectors that contain the nucleic acid sequence of human soluble guanylyl cyclase  $\alpha 1$  (hsGC $\alpha 1$ ) and/or human soluble guanylyl cyclase  $\beta 1$  (hsGC $\beta 1$ ) are used for prevention and therapy of the diseases mentioned above. Likewise, a mixture of two vectors in which one vector contains the nucleic acid sequence of human soluble guanylyl cyclase  $\alpha 1$  (hsGC $\alpha 1$ ) and the second that of human soluble guanylyl cyclase  $\beta 1$  (hsGC $\beta 1$ ) can be used for somatic gene transfer. Especially preferred is somatic gene transfer into endothelial cells, vascular smooth muscle cells, neointimal cells, fibroblasts, and other vascular cells as well as into blood particles (platelets, leukocytes, and others) and the liver.

The methods of gene transfer described in the present invention can also be used for gene transfer of human soluble guanylyl cyclase  $\alpha 2$  (GeneBank: x63282) and of the human homologue of soluble guanylyl cyclase  $\beta 2$  (from rat; GeneBank: m57507) as well as for other human soluble guanylyl cyclases.

Another object of the present invention relates to antibodies to human soluble guanylyl cyclase  $\alpha 1/\beta 1$  (hsGC $\alpha 1/\beta 1$ ), obtainable by immunization of a mammal with hsGC $\alpha 1/\beta 1$ , either of subunits  $\alpha 1$  or  $\beta 1$ , or immunogenic peptide fragments thereof, as well as isolation of the antibodies.

#### Figures:

Figure 1 shows various possibilities of modulation of soluble guanylyl cyclase (CsGC- $\alpha 1$ /sGC- $\beta 1$ ). Normal activation is mediated by NO synthase (NOS) and NO. NO, however, reacts with oxygen radicals to form peroxynitrite (ONOO $^-$ ), which is cytotoxic and only poorly activates sGC. NO can also be released by NO donors such as nitroglycerin or sodium nitroprusside. sGC can be directly activated by modulators of sGC (e.g. YC-1); alternatively, the activation by NO of sGC is potentiated by these modulators. Additionally, sGC can be overexpressed by use of gene transfer (e.g. using adenoviral vectors) or a pathologically low expression level of sGC can be compensated. Adenoviral (or other) vectors with mutated sGC that

has a higher basal activity can also be used. Thus, an elevated cGMP level could be achieved permanently independent of NOS, NO, NO donors, or sGC modulators.

Figure 2 shows a schematic comparison of the bovine and rat sGC $\alpha$ 1 subunits together with the published sequence of the human cDNA clone termed 'sGC $\alpha$ 3' (Giuli et al., 1992). The bars represent the protein. 'N' represents the N-terminus and 'C' the C-terminus. The functional segments 'regulatory domain', 'sGC homology domain', and 'cyclase domain' of these proteins are marked with different patterns. The regions 'S1' and 'S2' for which no homologous regions are found in the bovine and rat sGC $\alpha$ 1 proteins are marked in black.

Figure 3 shows a schematic illustration of the human sGC $\alpha$ 3 clone with the sequence errors published (Giuli et al., 1992). The cDNA is shown above; the bar represents the coding region of the cDNA, the lines left and right of the bar represent the 5' and 3' untranslated regions, respectively. 'S1' and 'S2' represent the regions that have no homology to the bovine and rat isoforms of sGC $\alpha$ 1 (see Fig. 2). The positions of the sequence errors are marked below: line a shows the nucleotide insertions, line b the deletion, and line c the exchanges. A base-pair (bp) scale is shown below. The sequence errors for each of the 3 lines a, b, and c are listed at the bottom: the letter specifies the type of the base concerned, and the number its position in the cDNA.

Figure 4 shows the verification of the expression of human sGC $\alpha$ 1 (A) and sGC $\beta$ 1 (B) in human tissues by means of PCR using cDNA libraries. Shown is a photo of an ethidium bromide-stained agarose gel under UV light with separated PCR products. The arrow on the left points to the specific product. The PCR primers are visible at the bottom of the photo. The tissues from which the cDNA libraries were produced are indicated at the top. No cDNA was added in the negative control, and in the positive control, plasmid containing the cDNA of hsGC $\alpha$ 1 was added.

Figures 5 and 6 show the baculovirus transfer vectors pVL1393 and pAcG2T, respectively (both without the hsGC cDNA), which were used for the construction of recombinant baculoviruses for the expression of human sGC $\alpha$ 1/ $\beta$ 1 in Sf9 cells. The circular plasmid with the restriction sites (short names and position in base pairs), the gene for ampicillin resistance (Amp<sup>R</sup>), the 'origin of replication' (ColE ori), the

polyhedrin promotor, the glutathione-S-transfease sequence (only in Fig. 6), and the multiple cloning site (MCS) are shown at the top of the figures. Figure 5 shows the multiple cloning site with its unique restriction sites at the bottom. Figure 6 shows the multiple cloning site with the unique restriction sites as well as a thrombin cleavage site normal at the bottom.

Figure 7 shows the construction of the plasmids hsGC $\beta$ 1-pVL1393 (without GST-tag) with the hsGC $\beta$ 1 cDNA, which was used to obtain genetically modified, hsGC $\beta$ 1-expressing baculovirus by homologous recombination. The procedure for the plasmid pAcG2T-hsGC $\beta$ 1 (with GST-tag = glutathione-S-transferase cDNA from *Schistosoma japonicum*) is identical. A fragment was produced that carries an additional BamHI-site at its 5' end by means of PCR with the primers A (bases 89-116 of the hsGC $\beta$ 1 cDNA + BamHI site at its 5' end) and B (bases 692-711 of the hsGC $\beta$ 1 cDNA [noncoding strand] with natural KpnI site). Due to the additional restriction sites, fragment 1 (PCR fragment with new BamHI site) and fragment 2 (hsGC $\beta$ 1 cDNA from KpnI site to EcoRI site) could be inserted together into the BamHI and EcoRI sites of the plasmid pVL1393. Thus, the complete hsGC $\beta$ 1 cDNA is under the control of the polyhedrin promotor (PHP).

Figure 8 shows the construction of the plasmid hsGC $\alpha$ 1-pVL1393 (without GST-tag) with the hsGC $\alpha$ 1 cDNA which was used to obtain genetically modified, hsGC $\alpha$ 1-expressing baculovirus by homologous recombination. The procedure for the plasmid pAcG2T-hsGC $\alpha$ 1 (with GST-tag = glutathione-S-transferase cDNA from *Schistosoma japonicum*) is identical. A fragment was produced that carries an additional BamHI site at its 5' end and a natural BsaAI site within the sequence by means of PCR with the primers C (bases 524-541 of the hsGC $\alpha$ 1 cDNA + BamHI site at its 5' end) and D (bases 1232-1249 of the hsGC $\alpha$ 1 cDNA [noncoding strand]). Due to the added restriction site, fragment 3 cut with BsaAI (PCR fragment with new BamHI site to the BsaAI site) and fragment 4 (hsGC $\alpha$ 1 cDNA from BsaAI site to EcoRI site) could be inserted together into the BamHI and EcoRI sites of the plasmid pVL1393. Thus, the complete hsGC $\alpha$ 1 cDNA is under the control of the polyhedrin promotor (PHP).

Figure 9 shows the verification of the expression of hsGC $\alpha$ 1/ $\beta$ 1 in Sf9 cells, which were infected with the genetically modified viruses described above (with hsGC $\alpha$ 1 or hsGC $\beta$ 1 cDNA; both without GST-tag = glutathione-S-transferase cDNA from *Schistosoma japonicum*). On the left (A) a Coomassie-stained 10% SDS-polyacrylamide gel is shown on which were loaded the cell homogenates that had been separated into pellet (P) and supernatant (S) by centrifugation (20,000 x g). 'Co' designates the control with noninfected Sf9 cells. ' $\alpha$ 1' designates Sf9 cells that were infected with viruses containing the hsGC $\alpha$ 1 cDNA, and ' $\beta$ 1' designates Sf9 cells that were infected with viruses containing the hsGC $\beta$ 1 cDNA. The position of hsGC $\alpha$ 1 and hsGC $\beta$ 1 in the gel are marked ( $\alpha$ 1 or  $\beta$ 1). On the right an immunoblot is shown with supernatant (S) and pellet (P) of the cell homogenate from Sf9 cells that were either noninfected (Co) or coinfecting with hsGC $\alpha$ 1 and hsGC $\beta$ 1 baculoviruses ( $\alpha$ 1+ $\beta$ 1). The peptide antibodies to hsGC $\beta$ 1 described above (anti-hsGC $\beta$ 1) were used in immunoblotting first (Fig. 9B, lines 1-4). Afterwards, the blot was redeveloped with the peptide antibodies to hsGC $\alpha$ 1 (anti-hsGC $\alpha$ 1; Fig. 9B, lanes 5-8), which additionally revealed the hsGC $\beta$ 1 bands.

Figure 10 shows the activity of guanylyl cyclase (formation of cGMP from GTP) in intact Sf9 cells that were coinfecting with the genetically modified baculoviruses containing hsGC $\alpha$ 1 and hsGC $\beta$ 1 cDNA described here (both without GST-tag). Shown is the content of cGMP in pmol per  $10^6$  cells with different treatments, indicated at the bottom of the figure. Sample 1 is untreated in both panels A and B. 1 mM IBMX (3-isobutyl-1-methylxanthine) was added to each of the other samples (top line: black crossbar). In the middle line, the concentration of SNP added to the samples is indicated in  $\mu$ M. The bottom line shows the concentration of added YC-1 (A, left) or ODQ (B, right) in  $\mu$ M.

YC-1 = 3-(5'-hydroxymethyl-2'-furyl)-1-benzylindazol;

ODQ = 1H-[1,2,4]oxadiazolo[4,3,-a]chinoxalin-1-on;

SNP = sodium nitroprusside;

cGMP = cyclic 3', 5'-guanosine monophosphate

Figure 11 shows the activity of guanylyl cyclase (formation of cGMP from GTP) in homogenates of Sf9 cells that were coinfecting with the genetically modified baculoviruses containing hsGC $\alpha$ 1 and hsGC $\beta$ 1 cDNA described here (both without

GST-tag). As shown at the top, (A), the soluble cell fraction (supernatant after centrifugation at 20,000 x g) was used, and at the bottom, (B), the respective pellet. In each case, the amount of cGMP formed in pmol per mg protein per minute is shown for homogenates of cells that were harvested at different time points (indicated in hours) after infection with the baculoviruses. cGMP formation was measured with (black boxes) and without (white boxes) the addition of 100  $\mu$ M SNP.

Figure 12 shows the activity of guanylyl cyclase (formation of cGMP from GTP) in Sf9 cells that were coinfectd with the genetically modified baculoviruses containing hsGC $\beta$ 1 cDNA (without GST-tag) and hsGC $\alpha$ 1 cDNA (with GST-tag = glutathione-S-transferase cDNA from *Schistosoma japonicum*). The formation of cGMP in pmol per mg protein per minute during the purification procedure (affinity chromatography on glutathione sepharose 4B) is shown. In each case, activity was measured in the lysate (after removing the insoluble part by centrifugation at 20,000 x g), in the supernatant after binding of hsGC to glutathione sepharose 4B (flow-through), in both supernatants of the washes of hsGC bound to glutathione sepharose 4B (1. and 2. wash), as well as in the supernatant after elution of hsGC with reduced glutathione (1. and 2. elution). The formation of cGMP was determined without (black boxes, 'basal') and with (grey boxes, '+ 100  $\mu$ M SNP') addition of 100  $\mu$ M SNP.

Figure 13 shows the natural, endogenous expression of hsGC $\alpha$ 1 and hsGC $\beta$ 1 in different human tissues in an immunoblot.

The peptide antibodies to hsGC $\alpha$ 1 described above (anti-hsGC $\alpha$ 1) were used on the left (A), and the peptide antibodies to hsGC $\beta$ 1 (anti-hsGC $\beta$ 1) were used on the right (B). On the right, the peptide to which the antibodies were raised was added as negative control (peptide: +), while no peptide was added on the left (peptide: -). The SDS extracts of rhsGC $\alpha$ 1- (in panel A of the figure) or rhsGC $\beta$ 1-overexpressing Sf9 cells (in panel B of the figure) (Sf9), of human cerebral cortex (cortex), of human cerebellum (cerebellum), and of human lung (lung) were loaded on an 8% polyacrylamide gel. The specific bands of hsGC $\alpha$ 1 ( $\alpha$ 1) and hsGC $\beta$ 1 ( $\beta$ 1) are indicated by an arrow.

Figure 14 shows in immunoblots the purification of hsGC $\alpha$ 1 (as a hsGC $\alpha$ 1/hsGC $\beta$ 1 dimer) from Sf9 cells that were coinfectd with the genetically modified baculoviruses

containing hsGC $\alpha$ 1 cDNA with GST-tag (= glutathione-S-transferase cDNA from *Schistosoma japonicum*) and hsGC $\beta$ 1 cDNA (without GST-tag). Cell lysate was incubated with glutathione sepharose 4B, and after binding, the supernatant was loaded (supernatant after binding). The sepharose was washed twice and the  
5    respective supernatants of this wash were loaded (1. and 2. wash). Subsequently, elution was performed by cleavage of the hsGC $\alpha$ 1 protein from the GST-tag with thrombin and an aliquot was loaded ('elution with thrombin'). Then, SDS stop buffer was added to the glutathione sepharose 4B and an aliquot was loaded (GSH sepharose after elution). In addition, glutathione sepharose 4B with bound hsGC $\alpha$ 1  
10    without prior thrombin elution to which SDS buffer had been added was loaded (GSH sepharose before elution). The immunoblot was developed with the affinity-purified peptide antibodies to the C-terminus of hsGC $\alpha$ 1 described here. The arrows on the right point to the specific bands of hsGC $\alpha$ 1 with the GST-tag (GST-hsGC $\alpha$ 1) and hsGC $\alpha$ 1 without the GST-tag (hsGC $\alpha$ 1).

15    Figure 15 shows the purification of hsGC $\alpha$ 1/ $\beta$ 1 from Sf9 cells that were coinfectd with the genetically modified baculoviruses containing hsGC $\alpha$ 1 cDNA with GST-tag (= glutathione-S-transferase cDNA from *Schistosoma japonicum*) and hsGC $\beta$ 1 cDNA without GST-tag in a Coomassie Brilliant Blue R250-stained SDS-polyacrylamide gel.  
20    Cell lysate of these infected Sf9 cells (lysate) was incubated with glutathione sepharose 4B, and the supernatant loaded after the binding (supernatant after binding). The glutathione sepharose 4B was washed twice and the respective supernatants of this wash buffer were loaded (1. and 2. wash). In one sample, the bound GST-hsGC $\alpha$ 1/ $\beta$ 1 was eluted by incubation with reduced glutathione and  
25    loaded (elution with glutathione). In the other samples, the glutathione sepharose was washed with the buffer for the thrombin cleavage—without thrombin—and the supernatant of this buffer was loaded (3. wash). Then, the hsGC $\alpha$ 1/ $\beta$ 1 dimer was eluted by incubation with different amounts of thrombin and the eluates were loaded (elution with 0.25-1 U/ml thrombin) [U = unit]. The same relative amount of each of  
30    the samples was used. The bands visible with the different elution methods are indicated on the right: GST-hsGC $\alpha$ 1 = hsGC $\alpha$ 1 with GST-tag; hsGC $\alpha$ 1 = hsGC $\alpha$ 1 without GST-tag; hsGC $\beta$ 1 = hsGC $\beta$ 1 without GST-tag. On the left are the molecular



weight standards loaded on the gel, the size of which (in kDa) is indicated on the far left side.

Figure 16 shows the construction of the recombinant adenoviral hsGC vectors. The cDNAs of hsGC $\alpha$ 1 and hsGC $\beta$ 1 (grey bars) were inserted in the adenoviral transfer plasmid pZS2, which has a deletion in the adenovirus E1 region ( $\Delta$ E1) and a unique XbaI site in this plasmid. This resulted in the plasmids hsGC $\alpha$ 1-pZS2 and hsGC $\beta$ 1-pZS2, respectively. hsGC $\alpha$ 1-pZS2 and hsGC $\beta$ 1-pZS2 cut with the restriction enzyme XbaI (middle bar, indicated as 'sGCpZS2') were ligated into the XbaI site of the long arm (upper bar, 'RR5') of Ad5. This resulted in the adenovectors Ad5CMVhsGC $\alpha$ 1 and Ad5CMVhsGC $\beta$ 1, respectively (lower bar, 'Ad 5 CMV sGC') in which the sGC cDNAs lie under the control of the CMV promotor and CMV enhancer (CMV = cytomegalovirus).

Figure 17 shows the stimulation of sGC activity by 100  $\mu$ M SNP (= sodium nitroprusside) in EA.hy926 cells that were coinfectd with both hsGC adenoviruses Ad5CMVhsGC $\alpha$ 1 and Ad5CMVhsGC $\beta$ 1 (samples A-C) and in noninfected EA.hy926 cells (sample D). The amount of pmol cGMP formed per mg protein per minute is plotted on the Y axis. The dark bars represent basal cGMP formation without SNP stimulation, and the light bars represent cGMP formation with SNP stimulation.

Figure 18 shows the DNA sequence of human soluble guanylyl cyclase  $\alpha$ 1 (hsGC $\alpha$ 1); SEQ ID NO: 1.

Figure 19 shows the amino acid sequence of human soluble guanylyl cyclase  $\alpha$ 1 (hsGC $\alpha$ 1); SEQ ID NO: 2.

Figure 20 shows the DNA sequence of human soluble guanylyl cyclase  $\beta$ 1 (hsGC $\beta$ 1); SEQ ID NO: 3.

Figure 21 shows the amino acid sequence of human soluble guanylyl cyclase  $\beta$ 1 (hsGC $\beta$ 1); SEQ ID NO: 4.

Figure 22 shows the amino acid sequence of the peptide that was used for the production of antibodies to human soluble guanylyl cyclase  $\alpha 1$  (hsGC $\alpha 1$ ) (corresponds to amino acids 634-647 of hsGC $\alpha 1$ ); SEQ ID NO: 5.

- 5 Figure 23 shows the amino acid sequence of the peptide that was used for the production of antibodies to human soluble guanylyl cyclase  $\beta 1$  (hsGC $\beta 1$ ) (corresponds to amino acids 593-614 of hsGC $\beta 1$ ); SEQ ID NO: 6.

- 10 Figure 24 shows the DNA sequence of the PCR primer pair for human soluble guanylyl cyclase  $\alpha 1$  (hsGC $\alpha 1$ ). Upper primer (corresponds to nucleotides 524-541 of the hsGC $\alpha 1$  cDNA sequence with added BamHI restriction site); SEQ ID NO: 7. Lower primer (corresponds to nucleotides 1249-1232 of the hsGC $\alpha 1$  cDNA sequence [noncoding strand]); SEQ ID NO: 8.

- 15 Figure 25 shows the DNA sequence of the PCR primer pair for human soluble guanylyl cyclase  $\beta 1$  (hsGC $\beta 1$ ). Upper primer (corresponds to nucleotides 89-106 of the hsGC $\beta 1$  cDNA sequence with added BamHI restriction site); SEQ ID NO: 9. Lower primer (corresponds to nucleotides 692-711 of the hsGC $\beta 1$  cDNA sequence [noncoding strand]); SEQ ID NO: 10.

20

Figure 26 shows the stimulation of sGC activity by 100  $\mu$ M SNP (sodium nitroprusside) in ECV 304 cells that had been coinfectd with different amounts of both hsGC adenoviruses Ad5CMVhsGC $\alpha 1$  and Ad5CMVhsGC $\beta 1$  (samples 1-6) and in noninfected ECV 304 cells (sample 7). The amount of pmol cGMP formed per mg protein per minute is plotted on the Y axis. The dark bars represent basal cGMP formation without SNP stimulation, and the light bars represent cGMP formation with SNP stimulation.

25

- 30 Figure 27 shows the stimulation of sGC activity by 100  $\mu$ M SNP (sodium nitroprusside) in A10 cells that had been coinfectd with different amounts of both hsGC adenoviruses Ad5CMVhsGC $\alpha 1$  and Ad5CMVhsGC $\beta 1$  (samples 1-6) and in noninfected A10 cells (sample 7). The amount of pmol cGMP formed per mg protein per minute is plotted on the Y axis. The dark bars represent basal cGMP formation

without SNP stimulation, and the light bars represent cGMP formation with SNP stimulation.

5 Figure 28 shows the stimulation of sGC activity by 100  $\mu$ M SNP (sodium nitroprusside) in ECV 304 cells that had been coinfectd with  $5 \times 10^{10}$  of each of the hsGC adenoviruses Ad5CMVhsGC $\alpha$ 1 and Ad5CMVhsGC $\beta$ 1 and then have been harvested at different time points as well as the activity in noninfected ECV 304 cells. The time point of the harvest, in days after infection (0, no infection), is plotted on the X axis. The amount of pmol cGMP formed per mg protein per minute is plotted on the  
10 Y axis. The dark bars represent basal cGMP formation without SNP stimulation, and the light bars represent cGMP formation with SNP stimulation.

Figure 29 shows the stimulation of sGC activity by 100  $\mu$ M SNP (sodium nitroprusside) in A10 cells that had been coinfectd with  $5 \times 10^{10}$  of each of the hsGC  
15 adenoviruses Ad5CMVhsGC $\alpha$ 1 and Ad5CMVhsGC $\beta$ 1 and then have been harvested at different time points as well as the activity in noninfected A10 cells. The time point of the harvest, in days after infection (0, no infection), is plotted on the X axis. The amount of pmol cGMP formed per mg protein per minute is plotted on the Y axis. The dark bars show basal cGMP formation without SNP stimulation, and the light bars  
20 represent cGMP formation with SNP stimulation.

Figure 30 shows the detection of sGC subunits in E304 cells and A10 cells, respectively, that had been coinfectd with  $5 \times 10^{10}$  of each of the hsGC  
25 adenoviruses Ad5CMVhsGC $\alpha$ 1 and Ad5CMVhsGC $\beta$ 1 for two days in a protein immunoblot with the peptide antibodies to hsGC $\alpha$ 1 and hsGC $\beta$ 1 described above. Lane 1, purified hsGC (according to Figure 12); lane 2, ECV 304 cells with hsGC gene transfer; lane 3, A10 cells with hsGC gene transfer; lane 4, purified hsGC (according to Figure 12).

Advantages of the present invention and solution of the above-mentioned technical problems according to the present invention

- 1.) The cDNA clones described in the literature as sGC $\alpha$ 3 and sGC $\beta$ 3 were identified as human homologues of bovine and rat sGC $\alpha$ 1 and sGC $\beta$ 1, and in the following are termed human sGC $\alpha$ 1 (hsGC $\alpha$ 1) and human sGC $\beta$ 1 (hsGC $\beta$ 1). According to present knowledge, this sGC $\alpha$ 1/ $\beta$ 1 isoform is pharmacologically more important due to its function in the cardiovascular system. Because the original clone of hsGC $\alpha$ 3 was examined, it could be shown that hsGC $\alpha$ 1 and hsGC $\alpha$ 3 do not exist in parallel but rather only the form hsGC $\alpha$ 1. Thus, an obvious target protein for pharmacological mass screening and gene therapy has been identified.
- 2.) With the methods described by the present invention, a functional, active expression of human sGC has been obtained for the first time. Thus, the respective protein can be produced by genetic methods for the first time.
- 3.) By use of the peptide antibodies to sGC of the present invention, it is possible to determine the expression of sGC in human tissues as well as diagnose dysfunctional conditions (if expression of sGC is too high, too low, or absent). In addition, the present invention provides the technical prerequisites needed to further elucidate the control of transcription and translation of hsGC. The peptide antibodies of the invention have the advantage that they are monospecific, directed at the human sequence, and that their suitability for immunoblots with human tissues has been demonstrated. Other peptide antibodies exhibit these features only partially: Harteneck et al. and Guthmann et al. used a peptide sequence (YKVVETVGDYMTVSGLP) that is relatively highly conserved in guanylyl cyclases. Thus, cross-reaction with particulate guanylyl cyclases (e.g. GC-C) would be expected. Furthermore, Guthmann et al. used a peptide sequence (YGPEVWEDIKKEA) identical to hsGC $\beta$ 1 and a peptide sequence identical to hsGC $\alpha$ 1 except for two amino acid exchanges (KKDVEEANANFLGKASGID), but the function of these antibodies in immunoblotting has only been shown for enriched hsGC from human platelets. In addition, these antibodies to hsGC $\alpha$ 1 recognized a second, unspecific product.

Humbert et al. and Koesling et al. used a peptide sequence (SRKNTGTEETEQDEN) from bovine sGC $\beta$ 1 that is in part (amino acids 1-10) identical to the peptide used here (amino acids 13-22) for hsGC $\beta$ 1, although the C-terminus (amino acids 11-15) differed markedly from that of the human sequence. The antiserum to this peptide, however, has not been tested on human protein and has only been used for immunoprecipitation of bovine sGC.

In addition to the peptides shown in Figures 22 and 23 and their immunogenic fragments for the production of the antibodies to hsGC $\alpha$ 1 or hsGC $\beta$ 1 in rabbits of the present invention, the production of monoclonal or polyclonal antibodies to the whole hsGC $\alpha$ 1/ $\beta$ 1 protein or its cleavage products is possible. Various animal species (preferably mouse, rat, or rabbit) can be used for the production of these antibodies.

4.) By use of the eukaryotic baculovirus/Sf9 expression system of the present invention, human soluble guanylyl cyclase  $\alpha$ 1/ $\beta$ 1 can be produced in high amounts. The attachment of a nucleotide sequence that codes for a polypeptide suitable for affinity chromatography (affinity tag, e.g. glutathione-S-transferase = GST-tag) with an attached protease cleavage site at the N-terminus of the cDNA of the  $\alpha$ 1 subunit allows the rapid and simple purification of the coexpressed dimeric protein by means of a single affinity chromatographic step. The attached affinity tag is subsequently removed by digestion with a protease. Thus, a protein identical in primary structure to the native protein is obtained. This revolutionary, fast, and clean purification yielding high amounts of very pure, functional human sGC results in new possibilities for a mass screening for specific activators and inhibitors as well as for the pharmacological characterization of potential drugs.

Purification of hsGC $\alpha$ 1/ $\beta$ 1 can also be accomplished by ion exchange chromatography, gel filtration, immunoaffinity chromatography, and other chromatographic procedures, e.g. on ATP-, GTP-, cGMP- or Blue-sepharose, and other similar chromatographic media.

5.) The process of the present invention can also be used for other isoforms of the human, rat, and bovine enzyme in an identical manner. In the process provided

in the present invention, various affinity tags (e.g. histidine oligomer) and different expression systems (e.g. *E. coli*) can be used. Other parts of the hsGC $\alpha$ 1 and hsGC $\beta$ 1 sequences can also be used for the production of antibodies to peptides or the whole protein.

5 6.) The availability of high amounts of an isolated human protein in high purity and in high quality is essential for modern pharmaceutical development. Until now, this requirement was not fulfilled in the search for alternatives to the classical NO donors. Mass screening for specific activators or inhibitors in animal testing is too  
10 expensive and would not make sense due to species differences, possible side effects, and effects on other isoforms. Cell culture systems have the disadvantage of requiring substantial additional effort to determine precisely at which point within the signaling cascades substances act. In addition, cell culture is expensive and touchy. Compared with the expression of a protein using  
15 recombinant DNA technology, the purification of a protein from animal tissues is more labor intensive and results in lower yields. In particular, the results of a pharmacological screening can only be generalized to a limited extent because of differences between species. In contrast, processes of the present invention provide for recombinant, inexpensive hsGC $\alpha$ 1/ $\beta$ 1 in high amounts, and  
20 unequivocal predictions concerning modulation, activation, or inhibition of this particular human enzyme can be made from a screening procedure. Species differences and lack of applicability to human are excluded a priori by the use of a human enzyme.

25 In addition, hsGC is available thereby in high amounts and adequate purity for crystallization and clarification of its structure. Thus, an important prerequisite for rational drug design by means of molecular modeling is fulfilled.

30 7.) In addition to the use of isolated hsGC $\alpha$ 1/ $\beta$ 1, intact Sf9 cells that express hsGC $\alpha$ 1/ $\beta$ 1 by being infected with the recombinant baculoviruses described here can be used in in vitro experiments.

8.) Transient (e.g. with adenoviral vectors) or stable overexpression can be achieved by means of gene transfer. Thus, the cGMP level can be elevated even at low

NO concentration or in the case of poor activation of sGC due to peroxynitrite formation. This approach also has conceptual advantages compared to the gene transfer of NOS (NO synthase) because the formation of cytotoxic peroxynitrite, which is less effective in activating sGC, is circumvented with the approach of the present invention. Furthermore, a permanent increase in the cGMP level could be achieved (e.g. for therapeutic purposes) independently of NOS, NO, NO donors, or sGC modulators by gene transfer of mutated sGC with elevated basal activity.

9.) In addition to the purification procedure for hsGC $\alpha$ 1/ $\beta$ 1 described here, purification from Sf9 cells is also possible after infection with the described baculoviruses containing hsGC $\alpha$ 1 or hsGC $\beta$ 1 cDNA if coinfection is performed with hsGC $\alpha$ 1 baculoviruses without a GST-tag (GST-tag = appended glutathione-S-transferase sequence from *Schistosoma japonicum*) and hsGC $\beta$ 1 baculoviruses having a GST-tag as well as with hsGC $\alpha$ 1 baculoviruses having a GST-tag and hsGC $\beta$ 1 baculoviruses having a GST-tag.

Furthermore, use of a column containing glutathione sepharose 4B is possible in addition to the use of glutathione sepharose 4B alone in a batch-wise step.

Purification can also be performed by digesting the dimeric fusion protein GST-hsGC $\alpha$ 1/ $\beta$ 1 with thrombin after elution of the fusion protein from the glutathione sepharose 4B with reduced glutathione. After dialysis (to remove the reduced glutathione), the GST-tag which has been cleaved from the protein can be removed from the mixture by additional affinity chromatography on glutathione sepharose 4B.

10.) The process of the present invention can also be used for human soluble guanylyl cyclase  $\alpha$ 2 (GeneBank: x63282) and any potentially existing human homologue of the soluble guanylyl cyclase  $\beta$ 2 (from rat; GeneBank: m57507) as well as for other human soluble guanylyl cyclases (in all technical variations described in the present invention).

11.) Using the adenoviral, somatic gene transfer of hsGC $\alpha$ 1 or hsGC $\beta$ 1 by means of coinfection with Ad5CMVhsGC $\alpha$ 1 or Ad5CMVhsGC $\beta$ 1, respectively, it was demonstrated in endothelial (ECV 304) and smooth muscle (A10) cells that the intracellular sGC activity measured as cGMP content after stimulation of sGC with SNP was increased from undetectable levels ( $\leq 100$  pmol/mg/min) to 12-fold higher values. An optimal concentration, regarding the amount of adenovirus (pfu) as well as an equimolar ratio of both adenoviruses (based on pfu), corresponding to the homodimeric structure of the enzyme, was determined. This effect on intracellular sGC activity was still detectable 15 days after a single infection, a period sufficient for the intended applications.

The examples illustrate the present invention.

#### Example 1

##### *The corrected sequence of hsGC $\alpha$ 1 and hsGC $\beta$ 1*

The original clone of the human isoform sGC $\alpha$ 3 and sGC $\beta$ 3 (Giulii et al., 1992) was sequenced again. While the sequence of the sGC $\beta$ 3 clone was confirmed (see SEQ ID NO: 3 and Figure 20), the sequencing of sGC $\alpha$ 3 showed that the original publication (Giulii et al., 1992) contained 19 sequencing errors, which are summarized in Figure 3. The corresponding corrected  $\alpha$ 3-cDNA sequence is shown in SEQ ID NO: 1 and Figure 18. The deduced amino acid sequence is shown in SEQ ID NO: 2 and Figure 19. Furthermore, the corrected sequence (see SEQ ID NO: 1 and Figure 18) is identical with the human sGC $\alpha$ 1 sequence published in the GeneBank (accession No. U58855), whereby the 5' untranslated region of the sequence provided here is 506 base pairs longer. Therefore, 'sGC $\alpha$ 3' is now classified as human sGC $\alpha$ 1 (hsGC $\alpha$ 1). Thus, it was shown that two different hsGC $\alpha$  subunits  $\alpha$ 1 and  $\alpha$ 3 do not both exist in humans, which could be important for cardiovascular research but rather—in analogy to the situation in bovine and rat tissues—only hsGC $\alpha$ 1.



**Table 1** Revised terminology of soluble guanylyl cyclase cDNAs and proteins and their detection in human tissue.

human sGC subunits		
	$\alpha$	$\beta$
isoform 1	cDNA and protein detectable	cDNA and protein detectable
	active, when coexpressed	
isoform 2	cDNA detectable	
	active, when coexpressed with bovine $\beta 1$	

- 5 The expression of sGC $\alpha 1$  and sGC $\beta 1$  mRNA in human tissues was demonstrated by means of PCR (Figure 4). The amplification of a hsGC $\beta 1$  fragment with a PCR primer pair (5'-AAAAGGATCCATGTACGGATTTGTGAAT-3' = nucleotides 89-106 of the hsGC $\beta 1$  cDNA sequence with added restriction site; 5'-ATGCGTGATTCTGGGTACC-3' = 692-711 of the hsGC $\beta 1$  cDNA sequence) with an annealing temperature of 54°C resulted in one specific band each in cDNA libraries from brain, heart, kidney, lung, pancreas, and skeletal muscle. The identity of the amplified fragment was confirmed by sequencing. The amplification of a hsGC $\alpha 1$  fragment with a PCR primer pair (5'-AAAAGGATCCATGTTCTGCACGAAGCTC-3' = nucleotide 524-541 of the hsGC $\alpha 1$  cDNA sequence with added restriction site; 5'-ATTATGGAAGCAGGAGG-3' = 1249-1232 of the hsGC $\alpha 1$  cDNA sequence) with an annealing temperature of 54°C resulted in one specific band each in cDNA libraries from heart (Figure 4A) and lung (not shown). In each case, the sequencing of the fragments resulted in the corrected hsGC $\alpha 1$  sequence; and the 'hsGC $\alpha 3$ ' sequence published by Giuli et al. was not found. Thus, it was shown that in humans, only one hsGC $\alpha 1/\beta 1$  exists, and that the potential hsGC $\alpha 3/\beta 3$  is a result of sequencing errors. This results in a clear picture for cardiovascular research concerning which sGC isoform should be the target protein for pharmacological screening.

### Example 2

#### *Construction of recombinant baculoviruses for the expression of human sGC $\alpha$ and sGC $\beta$ in insect cells*

- 5 In order to verify that hsGC $\alpha$ 1 and hsGC $\beta$ 1 can form a functional, heterodimeric sGC protein, both cDNAs were inserted into baculoviruses. Using these baculoviruses, the recombinant protein was expressed under the control of the strong polyhedrin promoter in insect cells (Sf9 cells). For the production of the recombinant baculoviruses, the baculovirus transfer vector pVL1393 (Pharmingen, San Diego, California, USA; Figure 5) and the baculovirus transfer vector pAcG2T (with glutathione-S-transferase sequence from *Schistosoma japonicum* and thrombin cleavage site; Pharmingen; Figure 6) were used in which the foreign genes hsGC $\alpha$ 1 and hsGC $\beta$ 1 were cloned, respectively. The cotransfection of such a recombinant pVL1393 or pAcG2T plasmid with BaculoGold baculovirus DNA (Pharmingen) allowed the direct isolation of the genetically modified baculoviruses with hsGC $\alpha$ 1 or hsGC $\beta$ 1 cDNA formed by homologous recombination from the cell culture media.

- The construction of pVL1393-hsGC $\beta$ 1 is schematically shown in Figure 7 (identical procedure for pAcG2T-hsGC $\beta$ 1). The coding region of the hsGC $\beta$ 1 cDNA with the 3' untranslated region but without the 5' untranslated region was cloned into pVL1393. For this, a BamHI site was introduced by means of PCR with primers A and B immediately upstream of the codon coding for the start methionine. The thus amplified fragment 1 was digested with BamHI/KpnI; fragment 2 was isolated from the sGC $\beta$ 1 cDNA clone with KpnI/EcoRI. Fragments 1 and 2 as well as the vector opened with BamHI and EcoRI were ligated (see Figure 7).

- The construction of pVL1393-hsGC $\alpha$ 1 is shown schematically in Figure 8 (identical procedure for pAcG2T-hsGC $\alpha$ 1). The coding region of the hsGC $\alpha$ 1 cDNA with the 3' untranslated region but without the 5' untranslated region was cloned into pVL1393. For this, a BamHI site was introduced by means of PCR with primers C and D immediately upstream of the codon coding for the start methionine. The thus amplified fragment 3 was digested with BamHI/BsaAI; fragment 4 was isolated from

the sGC $\alpha$ 1 cDNA clone with BsaAI/EcoRI. Fragments 3 and 4 as well as the vector opened with BamHI and EcoRI were ligated (see Figure 8).

For the production of the recombinant hsGC $\alpha$ 1 and hsGC $\beta$ 1 baculoviruses, the  
5 baculovirus transfer vectors (pVL1393-hsGC $\alpha$ 1, pAcG2T-hsGC $\alpha$ 1, pVL1393-hsGC $\beta$ 1, pAcG2T-hsGC $\beta$ 1) were each cotransfected with baculovirus DNA (Baculo-Gold; Pharmingen, San Diego, California, USA) in monolayers of Sf9 cell cultures. For this, the cells were cultivated at 27°C in IPL-41 media (Gibco) supplemented with 10% (vol/vol) fetal calf serum (Biochrom), 4% (vol/vol) tryptose-phosphate-broth  
10 (Gibco), 1% (vol/vol) Pluronic F68 (Gibco), 0,5% amphotericin B (Gibco), 80 µg/ml gentamycin sulfate (Gibco), and 0.5 mM  $\delta$ -aminolevulinic acid (Merck). Recombinant hsGC $\alpha$ 1 and hsGC $\beta$ 1 baculovirus clones were obtained from the culture media by means of plaque purification. For the production of virus stock solutions with high titer, Sf9 shaking cultures ( $0.5 \times 10^6$  cells/ml) were infected with a M.O.I. (multiplicity of infection) of 0.1 pfu/cell (pfu = plaque forming units) and harvested 6 days after infection.  
15

### Example 3

#### *Production of recombinant hsGC $\alpha$ 1 and hsGC $\beta$ 1 in Sf9 cells*

20 Ten of each recombinant hsGC $\alpha$ 1 and hsGC $\beta$ 1 baculovirus clones were tested for expression of recombinant protein in Sf9 cells. For this, Sf9 monolayer cell cultures were infected with plaque-purified, recombinant hsGC $\alpha$ 1 or hsGC $\beta$ 1 baculoviruses, incubated at 27°C for 5 days, harvested with a cell scraper, resuspended in 0.5 ml  
25 lysis buffer (25 mM TEA, pH 7.8, 50 mM NaCl, 1 mM EDTA, 5 mM DTT, 1 µM leupeptin, 0.5 mg/l trypsin inhibitor), and homogenized by sonication ('Sonifier 250', 'standard'-tip, Branson; 15 times, 'duty cycle': 15%, intensity: 1). After centrifugation of the homogenates at 20,000 x g, the supernatant and pellet were analyzed by SDS-PAGE. Three of the hsGC $\beta$ 1 baculovirus clones and two of the hsGC $\alpha$ 1 baculovirus  
30 clones yielded recombinant protein in amounts that were visible in the insoluble fraction by staining with Coomassie Brilliant Blue R250. Recombinant human sGC $\alpha$  and sGC $\beta$  (rhsGC $\alpha$  and rhsGC $\beta$ ) migrated with an apparent molecular weight of 79.5 (hsGC $\alpha$ 1) and 68.5 kDa (hsGC $\beta$ 1), which is very close to the predicted molecular

weights deduced from the amino acid sequence (77.5 and 70.5 kDa, respectively) (representative clones in Figure 9). The baculovirus clones that showed the highest expression of recombinant proteins in the immunoblot were used for the expression of functional heterodimeric hsGC [see examples 4-7].

5

#### Example 4

##### *Recombinant human sGC in intact insect cells is active and stimulated by NO*

- 10 For the production of functional heterodimeric human sGC, rhsGC $\alpha$ 1 and rhsGC $\beta$ 1 were coexpressed in Sf9 cells with recombinant baculoviruses. For this, Sf9 monolayer cultures [ $2.5 \times 10^6$  cells/dish,  $\varnothing$  90 mm; supplements see example 2] were coinfectd with a M.O.I. (multiplicity of infection) of 2 pfu/cell of each recombinant baculovirus (hsGC $\alpha$ 1 and hsGC $\beta$ 1; both without GST-tag) and cultivated for 48 hours
- 15 at 27°C. The basal as well as the NO-stimulated activity of the sGC in the cells was determined by measurement of the cGMP content in the cells in the presence of the phosphodiesterase inhibitor IBMX (3-isobutyl-1-methylxanthine).

- For the determination of the cGMP content, the culture media was replaced by
- 20 Krebs-Ringer-buffer (KRB; 119 mM NaCl; 4.74 mM KCl; 2.54 mM CaCl<sub>2</sub>; 1.19 mM MgSO<sub>4</sub>; 1.19 mM KH<sub>2</sub>PO<sub>4</sub>; 25 mM NaHCO<sub>3</sub>; 10 mM HEPES; pH 7.4; 0.1% BSA) which, in addition, contained 1 mM IBMX. The cells were incubated for one hour at 27°C. Then the cells were washed with ice-cold KRP and harvested in 1 ml ice-cold ethanol (80 %) with a cell scraper. The cells were homogenized by sonication [see
- 25 Example 3] and centrifuged at  $20,000 \times g$  for 20 minutes. The supernatant was dried in a speed-vac and the residue resuspended in 25 mM TEA, pH 7.8. The cGMP content was determined by means of RIA (Biotrend).

- The coexpression of rhsGC $\alpha$ 1 and rhsGC $\beta$ 1 resulted in the formation of functional
- 30 sGC with basal activity in Sf9 cells (Figure 10): While noninfected Sf9 cells contained approximately 0.1 pmol cGMP/ $10^6$  cells (not shown), approximately 20 pmol cGMP/ $10^6$  cells was found in rhsGC-expressing cells (Figure 10). This basal activity of recombinant hsGC was increased by an NO donor, SNP (sodium nitroprusside). When the cells were incubated with 10, 100, or 1000  $\mu$ M SNP for 2 minutes prior to

the harvest, the cGMP content increased in a concentration-dependent manner by up to 50 fold (Figure 10).

5

### Example 5

*Recombinant human sGC in insect cell extracts is active and stimulated by NO*

The activity of the recombinant hsGC (after expression with the recombinant baculoviruses described above) was not only determined in intact Sf9 cells but also in Sf9 cell extracts. For the production of such extracts, Sf9 shaking cultures [ $2 \times 10^6$  cells/ml; supplements see Example 2] were coinfectd with a M.O.I. (multiplicity of infection) of 1 pfu/cell of each virus (hsGC $\alpha$ 1 and hsGC $\beta$ 1; both without GST-tag) and incubated at 27°C. Samples were taken (4 ml) at 0, 24, 48, 72, 96, and 118 hours after infection, the cells sedimented, resuspended in 1 ml lysis buffer and homogenized by sonication [see Example 3]. The homogenates were centrifuged at 20,000 x g for 15 minutes and the insoluble pellet was resuspended again in lysis buffer. The samples were adjusted to 50% glycerin (vol/vol) and stored at -20°C. The protein concentration was determined spectrophotometrically with the standard method of Bradford (Bradford, 1976). The sGC activity was determined by the formation of [ $^{32}$ P]cGMP from [ $\alpha$ - $^{32}$ P]GTP (Schultz and Böhme, 1984). The reactions contained 50 mM TEA (pH 7.4), 3 mM MgCl<sub>2</sub>, 3 mM DTT, 1 mM IBMX, 1 mM cGMP, 5 mM creatine phosphate, 0.25 mg/ml creatine kinase, and 500  $\mu$ M GTP in a total volume of 100  $\mu$ l. After incubation at 37°C for 10 minutes, the reaction was started by simultaneous addition of cell extract and the sGC activators SNP, CO, or YC-1. The [ $^{32}$ P]cGMP formed was measured as described (Schultz and Böhme, 1984). Basal rhsGC activity (i.e. cGMP formation by rshGC *without* activation of the enzyme by addition of NO or other activators), mainly found in the soluble Sf9 cell fraction (Figure 11A), reached is maximum 72 hours after infection of the cells and was increased up to 5 fold with 100  $\mu$ M SNP (Figure 11A). The pellet fraction did not contain measurable basal sGC activity at any time point; although in the presence of SNP a low degree of sGC activity was found (Figure 11B).

### Example 6

#### *Influence of YC-1 and ODQ on recombinant human sGC*

YC-1 (3-(5'-hydroxymethyl-2'-furyl)-1-benzylindazol) and ODQ (1H-[1,2,4]oxadiazol[4,3,-a]chinoxalin-1-on) are substances that were described to specifically influence the activity of sGC. Thus, it was investigated if this also holds true for rhsGC.

After expression with the recombinant baculoviruses (without GST-tag) described above, rhsGC was activated in intact Sf9 cells by YC-1. The NO-potentiating effect was also found: the cGMP content of cells expressing rhsGC was increased 3.4 fold by incubation with 10  $\mu$ M YC-1 for 2 minutes (Figure 10A). 100  $\mu$ M YC-1 had the same effect (Figure 10A). When the cells were simultaneously treated with YC-1 and 100  $\mu$ M SNP, the cGMP levels doubled compared to the cGMP levels after stimulation with SNP alone (Figure 10A). Similar results were obtained with rhsGC in cell extracts.

ODQ is described as a selective inhibitor of NO-stimulated sGC that, however, does not inhibit the basal activity (Garthwaite et al., 1995). In Sf9 cells expressing rhsGC (after expression with the recombinant baculoviruses described above, without GST-tag), ODQ did not have an influence on the basal cGMP levels; the stimulation of rhsGC in intact cells with SNP, however, was inhibited by simultaneous incubation with ODQ (Figure 10B).

### Example 7

#### *Extraction of the purified human soluble guanylyl cyclase $\alpha 1/\beta 1$ (hsGC $\alpha 1/\beta 1$ )*

For the purification of recombinant human hsGC $\alpha 1/\beta 1$  from Sf9 cells, a recombinant baculovirus [see Example 2] was used in which a fusion protein composed of hsGC $\alpha 1$  and attached GST [so-called GST-tag, GST = glutathione-S-transferase from *Schistosoma japonicum*; see Example 2] is formed. Using this GST-tag, which binds to glutathione with high affinity, specific affinity chromatography on glutathione sepharose 4B (Pharmacia, Freiburg, Germany) can be performed. The Sf9 cells coinfectd with hsGC $\alpha 1$  baculoviruses (with GST-tag) and hsGC $\beta 1$  baculoviruses (without GST-tag) were lysed in 25 mM triethanolamine pH 7.8 / 1 mM EDTA / 5 mM DTT / 1  $\mu$ M leupeptin / 0.5  $\mu$ g/ml trypsin inhibitor / 0.2 mM PMSF (30 min hypotonic

lysis at 4°C). After addition of NaCl (75 mM final concentration), the homogenate was centrifuged at 75,000 x g for 1 hour at 4°C. The supernatant was mixed with the GSH sepharose 4B for 1 hour at room temperature. The glutathione sepharose 4B was then pelleted by centrifugation at 500 x g for 5 minutes and the supernatant removed.

- 5 A 10-fold volume of 50 mM Tris-HCl (pH 8.0) / 150 mM NaCl / 2.5 mM CaCl<sub>2</sub> / 0.1% 2-mercaptoethanol was added to the glutathione sepharose 4B and mixed for 1 minute. The mixture was centrifuged again at 500 x g for 5 minutes. The supernatant was removed and the glutathione sepharose 4B was washed again in the same way. To elute from the sepharose, the hsGCα1 protein (with the bound hsGCβ1) was
- 10 cleaved by thrombin from the GST-tag, which remained bound to the glutathione sepharose 4B at the specific cleavage site. Digestion with thrombin was performed in 50 mM Tris-HCl (pH 8.0) / 150 mM NaCl / 2.5 mM CaCl<sub>2</sub> / 0.1% 2-mercaptoethanol with 0.25 to 1 units thrombin/ml buffer for 1 or 3 hours at room temperature. The glutathione sepharose 4B (with the GST-tag) was pelleted again by centrifugation at
- 15 500 x g for 5 minutes and the supernatant containing the hsGCα1/β1 was removed. Another elution method was performed by addition of 50 mM Tris-HCl (pH 8.0) / 5 mM reduced glutathione and mixing for 30 minutes at room temperature. In this manner, hsGCα1/β1 with the GST-tag bound was removed from the glutathione sepharose 4B. After centrifugation at 500 x g for 5 minutes, the supernatant
- 20 containing dissolved GST-hsGCα1/β1 was removed.

By the elution with thrombin, a two-fold selectivity is achieved in a single affinity chromatographic step:

- 1.) Only proteins that have an affinity to reduced glutathione are able to bind.
- 25 2.) Of these proteins, only the proteins that are cleaved by thrombin will elute (as cleavage products).

The thrombin can be separated from the sample with a p-aminobenzamidine column to which thrombin binds specifically.

30

Figure 12 shows the specific enrichment of sGC activity after elution from the GSH sepharose 4B with glutathione compared with the activity in the lysate of infected Sf9 cells.

Figure 14 shows the binding of GST-hsGC $\alpha$ 1 to glutathione sepharose 4B and the cleavage of hsGC $\alpha$ 1 from the GST-tag by thrombin in an immunoblot.

Figure 15 shows the purification of coexpressed hsGC $\alpha$ 1 with GST-tag and hsGC $\beta$ 1 without GST-tag by affinity chromatography on glutathione sepharose 4B in a Coomassie Brilliant Blue R250-stained SDS-polyacrylamide gel. Upon elution with reduced glutathione, only two bands were visible, which correspond to GST-hsGC $\alpha$ 1 (larger product) and hsGC $\beta$ 1 (smaller product) (detected in an immunoblot). After elution with thrombin (0.25, 0.5, or 1 unit/ml for three hours at room temperature), however, the lower band was identical (hsGC $\beta$ 1; estimated molecular weight according to the migration in the gel of approximately 70 kDa), whereas the upper band was significantly smaller compared to elution with glutathione (hsGC $\alpha$ 1; estimated molecular weight of approximately 80 kDa) because the GST-tag was cleaved off by use of the thrombin elution. This approximately corresponds to the molecular weights of 77.5 kDa for hsGC $\alpha$ 1 and 70.5 kDa for hsGC $\beta$ 1 deduced from the amino acid sequences. In contrast to the elution with reduced glutathione, an additional very small band of approximately 25 kDa was visible after the thrombin elution, which is likely thrombin itself. Thrombin can be removed from the eluate by means of an aminobenzamidine sepharose column to which thrombin binds specifically. Additional bands were not detectable in this experiment.

### Example 8

#### *Production of polyclonal rabbit antisera to hsGC $\alpha$ 1 and hsGC $\beta$ 1*

Antisera were obtained by immunization of rabbits with synthetic peptides corresponding to sequences from hsGC $\alpha$ 1 (Phe-Thr-Pro-Arg-Ser-Arg-Glu-Glu-Leu-Pro-Pro-Asn-Phe-Pro [Figure 22/SEQ ID NO: 5]; amino acids 634-647) and from hsGC $\beta$ 1 (Lys-Gly-Lys-Lys-Glu-Pro-Met-Gln-Val-Trp-Phe-Leu-Ser-Arg-Lys-Asn-Thr-Gly-Thr-Glu-Glu-Thr [Figure 23/SEQ ID NO: 6] amino acids 593-614) that were coupled to KLH (keyhole limpet hemocyanin) via an additional C-terminal ( $\alpha$ 1) or N-terminal ( $\beta$ 1) cysteine residue. The antisera were affinity-purified with the corresponding peptides coupled to epoxy-activated sepharose (Pharmacia, Freiburg, Germany) according to the manufacturer's instructions.



### Example 9

#### *Detection of hsGC $\alpha$ 1 and hsGC $\beta$ 1 in different human tissues by immunoblotting*

- 5 Human lung tissue was obtained from a tumor-free area of a lung resection, and human cortex and cerebellum were from a normal autopsy. All tissues were immediately frozen in liquid nitrogen and stored at  $-70^{\circ}\text{C}$ . The frozen tissues were homogenized in a mortar and double-concentrated, hot SDS stop buffer (130 mM Tris-HCl, pH 6.8 / 16% [v/v] glycerol / 4% [w/v] SDS / 0.025% [w/v] bromophenol blue / 10 6.5% [v/v] 2-mercaptoethanol) was added to the powder. This was incubated at  $95^{\circ}\text{C}$  for 10 minutes and then centrifuged at  $20,000 \times g$  for 20 minutes. The supernatant was used for immunoblotting (for antibodies used, see above).

Expression of both subunits ( $\alpha$ 1 and  $\beta$ 1) was detected in all three tissues (Figure 13).

- 15 In contrast, expression could not be detected in kidney, liver, and pancreas (data not shown).

### Example 10

#### *Construction of recombinant adenoviral hsGC vectors*

20

- The cDNAs for hsGC $\alpha$ 1 and hsGC $\beta$ 1 were isolated from the original plasmid with the restriction enzyme EcoRI as 3.0 kb (hsGC $\alpha$ 1) and 2.4 kb (hsGC $\beta$ 3) fragments. The fragments were each inserted into the EcoRI restriction sites of the adenoviral transfer plasmid pZS2 (Figure 16), which contains an adenovirus type 5 sequence (Ad5) with a deletion in the E1 region ( $\Delta\text{E1}$ ), followed by an expression cassette with a CMV (cytomegalovirus) promoter/enhancer and a unique XbaI restriction site. 25 hsGC $\alpha$ 1-pZS2 and hsGC $\beta$ 1-pZS2 digested with XbaI were inserted into the XbaI site of the long arm (RR5) of Ad5 (Figure 16). The resulting recombinant adenoviral vectors Ad5CMVhsGC $\alpha$ 1 and Ad5CMVhsGC $\beta$ 1 are replication deficient because they lack the E1 region. To propagate the viruses, 293 cells that express E1 were 30 infected with these viruses. Viral plaques appeared 12-24 hours after the transfection. Viruses from single plaques were purified according to a standard procedure. Plaques containing recombinant viruses (Ad5CMVhsGC $\alpha$ 1 or Ad5CMVhsGC $\beta$ 1) were identified by means of PCR analysis: The plaque material

was freeze-thawed three times, incubated at 37°C for 30 minutes in lysis buffer (16.6 mM ammonium sulfate / 67 mM Tris-HCl pH 6.8 / 6.7 mM MgCl<sub>2</sub> / 5 mM 2-mercaptoethanol / 6.7 mM EDTA / 1.7 mM SDS / 50 µg/ml proteinase K), and thereafter heat inactivated for 10 minutes at 85°C. Finally, the DNA was isolated from the lysate with a standard phenol/chloroform extraction and used for PCR analysis.

#### Example 11

*Detection of cGMP formation in EA.hy926 cells after coinfection with the hsGC adenovectors Ad5CMVhsGCα1 and Ad5CMVhsGCβ1*

Ten 10-cm cell culture dishes with 'EA.hy926' cells were coinfectd with each of the adenovectors Ad5CMVhsGCα1 and Ad5CMVhsGCβ1 at  $2 \times 10^{10}$  pfu (plaque forming units) per dish. After 72 hours, the cells were harvested by adding hypotonic lysis buffer (25 mM triethanolamine pH 7.8 / 1 mM EDTA / 5 mM DTT / 1 µM leupeptin / 0.5 mg/l trypsin inhibitor / 0.2 mM PMSF) and detaching with a cell scraper. The homogenate was centrifuged at 500 x g for 15 minutes and the supernatant was mixed with an equal volume of glycerol and stored at -20°C. The stimulation of hsGC by 100 µM SNP (sodium nitroprusside) was determined by measurement of the basal cGMP level and the cGMP level after treatment with SNP according to the procedure described above (see Example 5). In three samples (A, B, C), a 7-fold to 10.75-fold elevation in the cGMP concentration compared to the basal activity was detectable after SNP stimulation, while no significant elevation was measurable in the control without adenovirus infection (Figure 17).

#### Example 12

*Detection of the cGMP formation in ECV 304 and A10 cells after coinfection with the hsGC adenoviruses Ad5CMVhsGCα1 and Ad5CMVhsGCβ1*

Three 10-cm cell culture dishes with 'ECV 304' cells were coinfectd with  $10^9 - 5 \times 10^{10}$  pfu (plaque forming units) of each of Ad5CMVhsGCα1 and Ad5CMVhsGCβ1. After 72 hours, the cells were harvested by two washes with PBS buffer (phosphate-buffered saline), addition of hypotonic lysis buffer, and then further treated according to the procedure described in Example 11. The samples were stored and the activity was determined (cGMP content in pmol/mg/min) as described in Example 5. A

significant basal activity (without SNP) and a significant maximal activity (with addition of 100  $\mu$ M SNP to the assay) was only detectable after the gene transfer of Ad5CMVhsGC $\alpha$ 1 and Ad5CMVhsGC $\beta$ 1 ( $5 \times 10^{10}$  pfu each) (Figure 26). In control experiments without Ad5CMVhsGC $\alpha$ 1 or Ad5CMVhsGC $\beta$ 1, no significant increase in cGMP formation by SNP was detectable. Qualitatively similar results were obtained in 'A10' vascular smooth muscle cells if these cells were also coinfecting with  $10^9 - 5 \times 10^{10}$  pfu Ad5CMVhsGC $\alpha$ 1 and Ad5CMVhsGC $\beta$ 1. Here, too, a significant basal activity (without SNP) and a significant maximal activity (with addition of 100  $\mu$ M SNP to the assay) was only detectable after the gene transfer of Ad5CMVhsGC $\alpha$ 1 and Ad5CMVhsGC $\beta$ 1 ( $5 \times 10^{10}$  pfu of each) (Figure 26). In control experiments without Ad5CMVhsGC $\alpha$ 1 or Ad5CMVhsGC $\beta$ 1, again no significant increase in cGMP formation by SNP was detectable (Figure 28).

#### Example 13

##### 15 *Kinetics of cGMP formation in ECV 304 and A10 cells after coinfection with the hsGC adenovectors Ad5CMVhsGC $\alpha$ 1 and Ad5CMVhsGC $\beta$ 1*

Three 10-cm cell culture dishes containing either 'ECV 304' or 'A10' cells were coinfecting with  $5 \times 10^{10}$  pfu (plaque forming units) Ad5CMVhsGC $\alpha$ 1 and Ad5CMVhsGC $\beta$ 1. The cells were harvested, and further treatment, storage, and determination of activity was performed as described in Examples 12, 11, and 5. Whereas at day 0 (before the coinfection), no significant increase in cGMP formation by SNP was detectable, 15 days after the gene transfer of Ad5CMVhsGC $\alpha$ 1 and Ad5CMVhsGC $\beta$ 1, a significant increase in cGMP formation after SNP addition was detectable (Figure 28). Qualitatively similar results were obtained in 'A10' vascular smooth muscle cells if these cells were also coinfecting with  $5 \times 10^{10}$  pfu Ad5CMVhsGC $\alpha$ 1 and Ad5CMVhsGC $\beta$ 1. Here, 15 days after coinfection, an increase in cGMP formation by SNP was also still detectable (Figure 29). Again, at day 0 (before the coinfection), there was still no significant increase in cGMP formation by SNP. In the same experiments, sGC subunits were detectable after adenoviral gene transfer, but not in noninfected control cells (Figure 30).

**References:**

- Bradford, M.M. (1976) *Annal. Biochem.* **72**: 248-254.
- 5 Friebe, A.; Schultz, G.; Koesling, D. (1996) *EMBO J.* **15**: 6863-6868.
- Garthwaite, J.; Southam, E.; Boulton, C. L.; Nielsen, E. B.; Schmidt, K.; Mayer, B. (1995) *Mol. Pharmacol.* **48**: 184-188.
- 10 Giuli, G.; Scholl, U.; Bulle, F.; Guellaen, G. (1992) *FEBS Lett.* **304**: 83-88.
- Guthmann, F.; Mayer, B.; Koesling, D.; Kukovetz, W.R.; Böhme, E. (1992) *Naunyn-Schmiedeberg's Arch. Pharmacol.* **346**: 537-541.
- 15 Harteneck, C.; Wedel, B.; Koesling, D.; Malkewitz, J.; Böhme, E.; Schultz, G. (1991) *FEBS Lett.* **292**: 217-222.
- Humbert, P.; Niroomand, F.; Fischer, G.; Mayer, B.; Koesling, D.; Hinsch, K.-D.; Gausepohl, H.; Frank, R.; Schultz, G.; Böhme, E. (1990) *Eur. J. Biochem.* **190**: 273-278.
- 20 Ko, F. N.; Wu, C. C.; Kuo, S.-C.; Lee, F.-Y.; Teng, C.M. (1994) *Blood* **84**: 4226-4233.
- Koesling, D.; Herz, J.; Gausepohl, H.; Niroomand, F.; Hinsch, K.-D.; Mülsch, A.; Böhme, E.; Schultz, G.; Frank, R. (1988) *FEBS Lett.* **239**: 29-34.
- 25 Schmidt, H. H. H. W.; Walter, U. (1994) *Cell* **78**: 919-925.
- Schultz, G.; Böhme, E. (1984) published in: Bergmeyer, H. U.; Bergmeyer, J.; Graßl, M. (Eds.): *Methods of Enzymatic Analysis*, Vol. 4, Verlag Chemie, Weinheim, pages 379-389.
- 30 Wu, C.-C.; Ko, F.-N.; Kuo, S.-C.; Lee, F.-Y.; Teng, C.-M. (1995) *British J. Pharmacol.* **116**: 1973-1978.

## Claims:

1. Isolated human soluble guanylyl cyclase  $\alpha 1$  (hsGC $\alpha 1$ ; SEQ ID NO: 2)/ $\beta 1$  (hsGC $\beta 1$ ; SEQ ID NO: 4) purified to apparent homogeneity.
2. A method for the production of  $\alpha 1$  (hsGC $\alpha 1$ ; SEQ ID NO:2) and  $\beta 1$  (hsGC $\beta 1$ ; SEQ ID NO:4) subunits of human soluble guanylyl cyclase comprising the expression in prokaryotic or eukaryotic host cells of expression vectors containing the DNA sequence of hsGC $\alpha 1$  and hsGC $\beta 1$  and obtaining the subunits.
3. The method for producing the  $\alpha 1$  and  $\beta 1$  subunits of human soluble guanylyl cyclase according to claim 2, wherein the step of obtaining the subunits comprises a lysis of the cells, the affinity chromatography of the cell lysate, and the subsequent elution of the subunits.
4. The method for producing the  $\alpha 1$  and  $\beta 1$  subunits of human soluble guanylyl cyclase according to claim 2 or 3, wherein the expression vector contains at least one additional DNA sequence coding for a domain for the specific affinity chromatography (affinity tag) with appended protease cleavage site.
5. The method for producing  $\alpha 1$  and  $\beta 1$  subunits of human soluble guanylyl cyclase according to claim 4, wherein the expression vector contains the DNA sequence for hsGC $\alpha 1$  with affinity tag, the DNA sequence for hsGC $\beta 1$  with affinity tag, the DNA sequence for hsGC $\alpha 1$  with affinity tag, and the DNA sequence for hsGC $\beta 1$ , the DNA sequence for hsGC $\beta 1$  with affinity tag and the DNA sequence for hsGC $\alpha 1$ , or the DNA sequence hsGC $\alpha 1$  with affinity tag and the DNA sequence for hsGC $\beta 1$  with affinity tag.
6. The method for producing human soluble guanylyl cyclase  $\alpha 1$  (hsGC $\alpha 1$ ; SEQ ID NO: 2)/ $\beta 1$  (hsGC $\beta 1$ ; SEQ ID NO: 4) comprising the separate expression in prokaryotic or eukaryotic host cells of an expression vector containing the DNA sequence for hsGC $\alpha 1$  or hsGC $\beta 1$ , extraction of the subunits, and

- reconstitution of subunits hsGC $\alpha$ 1 and hsGC $\beta$ 1 to form the dimeric guanylyl cyclase  $\alpha$ 1/ $\beta$ 1 (hsGC $\alpha$ 1/ $\beta$ 1).
7. The method for producing human soluble guanylyl cyclase  $\alpha$ 1/ $\beta$ 1 according to claim 6, wherein the step for the purification of the subunits consists of a separate lysis of cells containing hsGC $\alpha$ 1 or hsGC $\beta$ 1, the separate affinity chromatography of the cell lysates, and the subsequent elution of the subunits.
  8. The method for producing human soluble guanylyl cyclase  $\alpha$ 1 (hsGC $\alpha$ 1; SEQ ID NO: 2)/ $\beta$ 1 (hsGC $\beta$ 1; SEQ ID NO: 4) consisting of the coexpression of the DNA sequences of hsGC $\alpha$ 1 and hsGC $\beta$ 1 in prokaryotic or eukaryotic host cells, a lysis of the cells containing hsGC $\alpha$ 1 and hsGC $\beta$ 1, and affinity chromatography and subsequent elution of hsGC $\alpha$ 1/ $\beta$ 1.
  9. Use of a nucleotide sequence encoding the hsGC $\alpha$ 1 (SEQ ID NO: 2) and/or hsGC $\beta$ 1 (SEQ ID NO: 4) subunits of human soluble guanylyl cyclase  $\alpha$ 1/ $\beta$ 1 for somatic gene therapy.
  10. Use according to claim 9, wherein vectors or a mixture of vectors contain the nucleotide sequence of human soluble guanylyl cyclase  $\alpha$ 1 (hsGC $\alpha$ 1) and/or human soluble guanylyl cyclase  $\beta$ 1 (hsGC $\beta$ 1).
  11. Use according to claim 9 or 10 for the prevention and therapy of atherosclerosis and its complications, of restenosis, ischemia (infarction), peripheral arterial occlusive diseases, and arterial hypertension as well as for the prevention of atherosclerosis in patients with risk factors, transient ischemic attacks, cerebral ischemia, stroke (Apoplex), coronary heart disease, status post coronary bypass grafting, carotid stenosis, heart insufficiency and liver dysfunction, and as a supplement to therapy with sGC activators, sGC-sensitizing substances, NO donors, or phosphodiesterase inhibitors.
  12. Use according to claims 9 to 11, wherein the somatic gene transfer is carried out with endothelial cells, vascular smooth muscle cells, neointimal cells,

fibroblasts, or other vascular cells or blood particles (platelets, leukocytes, and others), or liver.

13. Antibodies against human soluble guanylyl cyclase  $\alpha 1$  (hsGC $\alpha 1$ ; SEQ ID NO: 2)/ $\beta 1$  (hsGC $\beta 1$ ; SEQ ID NO: 4) obtainable by immunization of a mammal with the peptide fragment Phe-Thr-Pro-Arg-Ser-Arg-Glu-Glu-Leu-Pro-Pro-Asn-Phe-Pro, or parts thereof, or immunogenic peptide fragments that overlap with this fragment, or obtainable by immunization of a mammal with the peptide fragment Lys-Gly-Lys-Lys-Glu-Pro-Met-Gln-Val-Trp-Phe-Leu-Ser-Arg-Lys-Asn-Thr-Gly-Thr-Glu-Glu-Thr or immunogenic fragment or immunogenic peptide fragments that overlap with this fragment and isolation of the antibodies.

**Abstract**

5 The invention relates to the expression of the cDNA clone for sub-units a1 (hsGCa1) and b1 (hsGCa1) of human soluble guanylylcyclase and the subsequent purification of the active enzyme and use thereof, the medical application of the expression of this clone by gene transfer, in addition to antibodies against peptides derived from said sequence and the use thereof.



Fig. 1






Figure 2



**S1**

**S2**

Figure 3

**a: nucleotide insertions**

C95, C367, T891, G900, T903, G913, T1006, G1074, G1487, A1488, A1489, G2108, G2555, T2560

**b: nucleotide deletions**


T between G111 and T112, T between T2128 and G2129, T between G2975 and T2976

**c: nucleotide exchanges**


C1525>G, G2125>A

Figure 4

A


PCR  
determination  
of hsGC $\alpha$ 1

B


PCR  
determination  
of hsGC $\beta$ 1


## Figure 5

### pVL1393 Baculovirus Transfer Vector



multiple cloning site (MCS) of pVL1393 with the unique restriction sites



**Figure 6****pAcG2T Baculovirus Transfer Vector**

multiple cloning site (MCS) of pAcG2T downstream of glutathione-S-transferase sequence (GST) with the thrombin cleavage site and the unique restriction sites






Figure 7: Cloning of hsGC $\beta$  in pVL1393

Primer: A 5' AAAA GGATCC ATGTACGGATTGTGAAT 3'  
BamHI (89) (116)

B 3' CCATGG GTCCTTAGTGCCTA 5'  
(692) KpnI (711)


Figure 8: Cloning of hsGC $\alpha$  in pVL1393

Primer: C 5' AAAA GGATCC ATGTTCTGCACGAAGCTC 3'  
BamHI (524) (541)

D 3' GGAGGGACGAAGGTATTA 5'  
(1232) (1249)




Figure 9



10/28

Fig. 10



A




Figure 12: Purification of GST-hsGCalpha1/beta1 on GSH-Sepharose  
4B




Figure 13





Figure 14



**Figure 15: Purification of hsGC  $\alpha$ 1/ $\beta$ 1 in a Coomassie stained SDS polyacrylamide gel**




Figure 16: Construction of the hsGC-adenovectors






Figure 17:  
Expression of human sGC in  
adenovirus-infected EA.hy926  
cells




Figure 18

CCCTTATGGC GATTGGGCGG CTGCAGAGAC CAGGACTCAG TTCCCTGCC CTAGTCTGAG  
 CCTACTGGGT GGGACTCAGC TCAGAGTCAG TTTTCCAGAA GCAGGTTTCA GTGCAGAGTT  
 TTCTCTACACT TTTCTCTGCC TAGAGCAGCG AGCAGCCTGG AACAGACCCA GGGCGAGGAC  
 ACCTGTGGGG GAGGGAGCGC CTGGAGCAGC TTAGAGACCC CAGCCGGGGC TGATCTCACC  
 ATGTGCGGAT TTGGGAGGCG CGCCCTGGAG CTGCTAGAGA TCCGGAAGCA CAGCCCCGAG  
 GTGTGCGAAG CCACCAAGAC TCGGGCTCTT GGAGAAAGCG TGAGCAGGGG GCCACCCGGG  
 TCTCCGGGCC TGTCTGCACC CTGTCCGCTG AGCTGCCTGA CAGTGACAAT GACATCCGAG  
 TTACCAGTGT CTTTGAATTG ATAGTGGCTT CTGTTTGTCG GTCTCATATA AGAATACAG  
 CTCATCAGGA GGAGATCGCA GCAGGGTAAG AGACACCAAC ACCATGTCTT GCACGAAGCT  
 CAAGGATCTC AAGATCACAG GAGAGTGCC TTTCTCCTTA CTGGCACGAG CTGGAGTTCC  
 TAACGAGTCT TCAGAGGAGG CAGCAGGAGG CTCAGAGAGC TGCAAGCAAA CCGTGCCCAT  
 CTGTCAAGAC ATTCTTGAGA AGAACATACA AGAAAGTCTT CCTCAAAGAA AAACCAAGTC  
 GAGCCGAGCT TATCTTCACA CTTTGGCAGA GAGTATTGCG AAAGTGAATT TCCCAGAGTT  
 TGAACGGCTG AATGTTGCAC TTCAGAGAAC ATTGGCAAAG CACAAAATAA AAGAAAGCAG  
 GAAATCTTTG GAAAGAGAAG ACTTTGAAAA AACAAATGCA GAGCAAGCAG TTTCAGCAGG  
 AGTTCCAGTG GAGGTATACA AAGAATCTCT TGGTGAAGAG GTTTTTAAAA TATGTTACGA  
 GGAATGAAAT AACATCCTTG GGGTGGTTGG AGSCACCTTT AAAGATTTTT TAAACAGCTT  
 CAGTACCCTT CTGAAACAGA GCAGCCATTG CCAAGAAGCA GGAAGAAAGG GCGAGCTTGA  
 GGACGCTCC ATTCTATGCC TGGATAAGGA GGATGATTTT CTACATGTTT ACTACTTCTT  
 CCTTAAGAGA ACCACCTCCC TGATTTCTCC CGGCATCATA AAGGCAGCTG CTCACCTATT  
 ATATGAAACG GAACTGGAAG TGTCTGTAAT GCCTCCCTGC TTCCATAAAT ATTGCAGCGA  
 GTTTGTGAAT CAGCCCTACT TGTGTACTC CGTTCACATG AAAAGCACCA AGCCATCCCT  
 GTCCCCAGC AAACCCAGT CCTCGCTGGT GATTTCCACA TCCTATTTCT GCAAGACATT  
 TCCATTCAT TTCTGTTTG ACAAAGATAT GACAATTTCTG CAATTGGGCA ATGGCATCAG  
 AAGGCTGATG AACAGGAGAG ACTTTCAAGG AAAGCCTAAT TTTGAAGAAT ACTTTGAAAT  
 TCTGACTCCA AAAATCAACC AGACGTTTAG CGGATCATG ACTATGTTGA ATATGCAGTT  
 TGTGTACGA GTGAGGAGAT GGGACAACCT TGTGAAGAAA TCTTCAAGGG TTATGGAACCT  
 CAAAGGCCAA ATGATCTACA TTGTTGAATC CAGTGCAATC TTGTTTTTGG GGTCAACCTG  
 TGTGGACAGA TTAGAAGATT TTACAGGAGC AGGGCTCTAC CTCTCAGACA TCCCAATTCA  
 CAATGCAGTG AGGAGATGTG TCTTAATAGG GGAACAAGCC CGAGCTCAAG ATGGCCTGAA  
 GAAGAGGCTG GGGAGCTGA AGGCTACCTT TGAGCAAGCC CACCAACGCC TGGAGGAGGA  
 GAAGAAAAAG ACAGTAGACC TTTCTGTGCTC CATATTTCCC TGTGAGGTTG CTCAGCAGCT  
 GTGGCAAGGG CAAGTTGTGC AAGCCAAGAA GTTCACTAAT GTCACCATGC TCTTCTCAGA  
 CATCGTTGGG TTCACTGCCA TCTGCTCCA GTGCTCAGG CTGCAGTGCA TCACCATGCT  
 CAATGCAGTG TACACTCGCT TCGACCAGCA GTGTGGAGAG CTGGATGTCT ACAAGGTGGA  
 GACCATTGGC GATGCTTATT GTGTAGCTGG GGGATTACAC AAAGAGAGTG ATAGCTCATG  
 TGTTCAGATA CGCTGATGG CCCTGAAGAT GATGGAAGCTC TCTGATGAAG TTATGTTCTC  
 CCGTAGGAAA CCTATCAAGA TCGGAATTGG ACTGCACTCT GGATCAGTTT TTGCTGGGCT  
 CGTTGGAGTT AAAATGCCCC GTTACTGTCT TTTTGGAAAC AATGTCACTC TGGCTAACAA  
 ATTTGAGTCC TGCAGTGTAC CACGAAAAAT CAATGTCAAG CATGTCAAGG CCAACAACCT  
 CAAAGACTGT CTTGTTTCG TGTPTACCCC TCGATCAAGG GAGGAACCTT CACCAAACTT  
 CCCTAGTGAA ATCCCCGGAA TCTGCCATT TCTGGATGCT TACCAACAGT GAACAACTC  
 AAAACACTGC TTCCAAAAGA AAGATGTGGA AGATGGCAAT GCCAATTTTT TAGGCAAAAG  
 ATCAGGAATA GATTAGCAAC CTATATACTT ATTTATAAGT CTTTGGGGTT TGACTCATTTG  
 AAGATTGTGA GAGCCTCTGA AAGCACTTTA GGGATTGTAG ATGGCTAACG AGCAGTATTA  
 AAAATTTGAG AGCCAAGTCA CAATCTTTCT CCGTTTAAAC ATGACAAAAT GTACTCACTT  
 CAGTACTTCA GCTCTTCAAG AAAAAAATA AAACCTTAAA TTAGCAACCT TGTACATATA  
 TTTCTATTAT ATAACCAGCA CTACTACCT GTACTCAAA TTAGCAACCT TGTACATATA  
 TCAGATAATT GTAGTCAATT GTACAAACTG ATGGAGTCAC CTGCAATCTC ATATCTCTGT  
 GGAATGCCAT GGTATTAAAT GTGTGTTTGT GATAGTGTG TCACAAAAAA AAAAAAATAA  
 AAAAAAATAA AAAAA

19/28

Figure 19

Met Phe Cys Thr Lys Leu Lys Asp Leu Lys Ile Thr Gly Glu Cys Pro  
 Phe Ser Leu Leu Ala Pro Gly Gln Val Pro Asn Glu Ser Ser Glu Glu  
 Ala Ala Gly Ser Ser Glu Ser Cys Lys Ala Thr Val Pro Ile Cys Gln  
 Asp Ile Pro Glu Lys Asn Ile Gln Glu Ser Leu Pro Gln Arg Lys Thr  
 Ser Arg Ser Arg Val Tyr Leu His Thr Leu Ala Glu Ser Ile Cys Lys  
 Leu Ile Phe Pro Glu Phe Glu Arg Leu Asn Val Ala Leu Gln Arg Thr  
 Leu Ala Lys His Lys Ile Lys Glu Ser Arg Lys Ser Leu Glu Arg Glu  
 Asp Phe Glu Lys Thr Ile Ala Glu Gln Ala Val Ala Ala Gly Val Pro  
 Val Glu Val Ile Lys Glu Ser Leu Gly Glu Glu Val Phe Lys Ile Cys  
 Tyr Glu Glu Asp Glu Asn Ile Leu Gly Val Val Gly Gly Thr Leu Lys  
 Asp Phe Leu Asn Ser Phe Ser Thr Leu Leu Lys Gln Ser Ser His Cys  
 Gln Glu Ala Gly Lys Arg Gly Arg Leu Glu Asp Ala Ser Ile Leu Cys  
 Leu Asp Lys Glu Asp Asp Phe Leu His Val Tyr Tyr Phe Phe Pro Lys  
 Arg Thr Thr Ser Leu Ile Leu Pro Gly Ile Ile Lys Ala Ala Ala His  
 Val Leu Tyr Glu Thr Glu Val Glu Val Ser Leu Met Pro Pro Cys Phe  
 His Asn Asp Cys Ser Glu Phe Val Asn Gln Pro Tyr Leu Leu Tyr Ser  
 Val His Met Lys Ser Thr Lys Pro Ser Leu Ser Pro Ser Lys Pro Gln  
 Ser Ser Leu Val Ile Pro Thr Ser Leu Phe Cys Lys Thr Phe Pro Phe  
 His Phe Met Phe Asp Lys Asp Met Thr Ile Leu Gln Phe Gly Asn Gly  
 Ile Arg Arg Leu Met Asn Arg Arg Asp Phe Gln Gly Lys Pro Asn Phe  
 Glu Glu Tyr Phe Glu Ile Leu Thr Pro Lys Ile Asn Gln Thr Phe Ser  
 Gly Ile Met Thr Met Leu Asn Met Gln Phe Val Val Arg Val Arg Arg  
 Trp Asp Asn Ser Val Lys Lys Ser Ser Arg Val Met Asp Leu Lys Gly  
 Gln Met Ile Tyr Ile Val Glu Ser Ser Ala Ile Leu Phe Leu Gly Ser  
 Pro Cys Val Asp Arg Leu Glu Asp Phe Thr Gly Arg Gly Leu Tyr Leu  
 Ser Asp Ile Pro Ile His Asn Ala Leu Arg Asp Val Val Leu Ile Gly  
 Glu Gln Ala Arg Ala Gln Asp Gly Leu Lys Lys Arg Leu Gly Lys Leu  
 Lys Ala Thr Leu Glu Gln Ala His Gln Ala Leu Glu Glu Lys Lys  
 Lys Thr Val Asp Leu Leu Cys Ser Ile Phe Pro Cys Glu Val Ala Gln  
 Gln Leu Trp Gln Gly Gln Val Val Gln Ala Lys Lys Phe Ser Asn Val  
 Thr Met Leu Phe Ser Asp Ile Val Gly Phe Thr Ala Ile Cys Ser Gln  
 Cys Ser Pro Leu Gln Val Ile Thr Met Leu Asn Ala Leu Tyr Thr Arg  
 Phe Asp Gln Gln Cys Gly Glu Leu Asp Val Tyr Lys Val Glu Thr Ile  
 Gly Asp Ala Tyr Cys Val Ala Gly Gly Leu His Lys Glu Ser Asp Thr  
 His Ala Val Gln Ile Ala Leu Met Ala Leu Lys Met Met Glu Leu Ser  
 Asp Glu Val Met Ser Pro His Gly Glu Pro Ile Lys Met Arg Ile Gly  
 Leu His Ser Gly Ser Val Phe Ala Gly Val Val Gly Val Lys Met Pro  
 Arg Tyr Cys Leu Phe Gly Asn Asn Val Thr Leu Ala Asn Lys Phe Glu  
 Ser Cys Ser Val Pro Arg Lys Ile Asn Val Ser Pro Thr Thr Tyr Arg  
 Leu Leu Lys Asp Cys Pro Gly Phe Val Phe Thr Pro Arg Ser Arg Glu  
 Glu Leu Pro Pro Asn Phe Pro Ser Glu Ile Pro Gly Ile Cys His Phe  
 Leu Asp Ala Tyr Gln Gln Gly Thr Asn Ser Lys Pro Cys Phe Gln Lys  
 Lys Asp Val Glu Asp Gly Asn Ala Asn Phe Leu Gly Lys Ala Ser Gly  
 Ile Asp End

Figure 20

```

CCCCCCCCG CCGCTGCCG CTCTGCGCTGG GTCCCTTCGG CCGTACCTCT CCGTGGGGGG
TGCCCTCCCCG GCTCCCGGTG CAGACACCAT GTACGGATT GTGAATCAG CCCTGGAGTT
GCTGGTGATC CGCAATTACG GCCCCGAGGT GTGGGAAGAC ATCAAAAAAG AGGCACAGTT
AGATGAAGAA GGCACAGTTTCT TTGTCAGAAT AATATATGAT GACTCCAAAT CTTATGATTT
GGTTGCTGCT CCAAGCAAAG TCCTCAATCT CAATGCTGGA GAAATCCTCC AAATGTTTGG
GAAGATGTTT TTGCTCTTTT GCCAAGAATC TGTTATGAT ACAATCTTC GTGCTCTGGG
CTCTAAATGTC AGAGAAATTC TACAGAACCT TGATGCTCTG CAGCACCACC TTGCTACCAT
CTACCCAGGA ATCGCTGCAC CTTCCTTTAG GTGCACTGAT GCAGAAAAAG GCAAGGACT
CATTTTGCAC TACTACTCAG AGAGAGAAGG ACTTCAGGAT ATTGTCATTG GAATCATCAA
AACAGTGGCA CAACAAATCC ATGGCACTGA AATAGACATG AAGGTTATTG AGCAAGAAAA
TGAAGAAATG TATCATACTC AATTTTTAAT TGAAGAAAA GAGTCAAAAG AAGAGGATTT
TTATGAAGAT CTGACAGAT TTGAAGAAAA TGGTACCAG GAATCAGCA TCAGCCCAT
TACATTCTCG AAAGCTTTTC CTTTTCATAT AATATTGAC CGGGACCTAG TGGTCACTCA
GTGTGGCAAT GCTATATACA GAGTTCTCCC CCAGCTCCAG CCTGGGAATT GCAGCCTTCT
GTCTGTCTTC TCGTGGTTTC GTCCTCATAT TGATATTAGT TTCCATGGGA TCCTTTCTCA
CATCAATACT GTTTTGTAT TGAGAAGCAA GGAAGGATTG TTGGATGTGG AGAAATTAGA
ATGTGAGGAT GAAGTGAATG GGACTGAGAT CAGCTGCTTA CGTCTCAAGG CTCAAATGAT
CTACTTACCT GAAGCAGATA GCATACTTTT TCTATGTCCA CCAAGTGCTA TGAACCTCGA
CGAATTGACA AGGAGAGGGC TGTATCTAAG TGACATCCCT CTGCATGAT CCACGCCGGA
TCTTGTTCCT TTGGGAGAAC AATTTAGAGA GGAATACAAA CTCACCAAG AACTGGAAAT
CCTCACTGAC AGGCTACAGC TCACGTTAAG AGCCCTGGAA GATGAAAGA AAAAGACAGA
CACATTGCTG TATTCTGTCC TTCTCCGCT TGTGCCAAT GAGCTGCGGC ACAAGCGTCC
AGTGCCTGCC AAAAGATATG ACAATGTGAC CATCTCTTT AGTGGCATTG TGGGCTTCAA
TGCTTTCTGT AGCAAGCATG CATCTGGAGA AGGAGCCATG AAGATCGTCA ACCTCTCAA
GCAGCTCTAC ACCAGATTG ACACACTGAC TGTATCCGGG AAAAACCCAT TTGTTATATA
GGTGGAGACT GTTGGTGACA ACTATATGAC AGTGAGTGGT TTACCAGAGC CATGCATTCA
CCATGCACGA TCCATCTGCC ACCTGGCCTT GGACATGATG GAAATGCTG GCCAGGTTCA
AGTAGATGGT GAATCTGTTC AGATAACAAT AGGGATACAC ACTGGAGAGG TAGTTACAGG
TGTCATAGGA CAGCGGATGC CTCGATACTG TCTTTTGGG AATACTGTCA ACCTCACAAG
CCGAACAGAA ACCACAGGAG AAAAGGGAAA AATAAATGT TCTGAATATA CATACAGATG
TCTTATGTCT CCAGAAATTT CAGATCCACA ATTCCACTTG GAGCACAGAG GCCAGTGT
CATGAAGGGC AAAAAGAAC CAATGCAAGT TTGGTTTCTA TCCAGAAAA ATACAGGAAC
AGAGGAACA AAGCAGGATG ATGACTGAAT CTGGATTAT GGGGTGAAGA GGAGTACAGA
CTAGTTTCCA GTTTCTCTCT AACACGTGCC AAGCCAGGA GCATTTCTTC CCTATGGATA
CAGATTTTCT TTGTCTCTTG TCCATTACCC CAAGACTTTC TTCTAGATAT ATCTCTCACT
ATCCGTTATT CAACCTTAGC TCTGCTTCT ATTACTTTTT AGGCTTAGT ATATTATCTA
TATCTTGGCT TTTGATGTGG ATGATGTGAG CTTCATGTGT CTTAAATCT ACTACAAGCA
AAAGCTAACA TGGTGAATCT CAAGTAGTAG GCACCCAATA AATATTGTT GAATTTAGTT
AAATGAAACT GAACAGTGT TGGCCATGTG TATATTTATA TCATGTTTAC CAAATCTGTT
TAGTGTCCA CATATATGTA TATGTATATT TTAATGACTA TAATGTAATA AGTTTATAT
CATGTTGGTG TATATCATT TAGAATCAT TTTCTAAAGG AGT

```

Figure 21

Met Tyr Gly Phe Val Asn His Ala Leu Glu Leu Leu Val Ile Arg Asn  
 Tyr Gly Pro Glu Val Trp Glu Asp Ile Lys Lys Glu Ala Gln Leu Asp  
 Glu Glu Gly Gln Phe Leu Val Arg Ile Ile Tyr Asp Asp Ser Lys Thr  
 Tyr Asp Leu Val Ala Ala Ala Ser Lys Val Leu Asn Leu Asn Ala Gly  
 Glu Ile Leu Gln Met Phe Gly Lys Met Phe Phe Val Phe Cys Gln Glu  
 Ser Gly Tyr Asp Thr Ile Leu Arg Val Leu Gly Ser Asn Val Arg Glu  
 Phe Leu Gln Asn Leu Asp Ala Leu His Asp His Leu Ala Thr Ile Tyr  
 Pro Gly Met Arg Ala Pro Ser Phe Arg Cys Thr Asp Ala Glu Lys Gly  
 Lys Gly Leu Ile Leu His Tyr Tyr Ser Glu Arg Glu Gly Leu Gln Asp  
 Ile Val Ile Gly Ile Ile Lys Thr Val Ala Gln Gln Ile His Gly Thr  
 Glu Ile Asp Met Lys Val Ile Gln Gln Arg Asn Glu Glu Cys Asp His  
 Thr Gln Phe Leu Ile Glu Glu Lys Glu Ser Lys Glu Glu Asp Phe Tyr  
 Glu Asp Leu Asp Arg Phe Glu Glu Asn Gly Thr Gln Glu Ser Arg Ile  
 Ser Pro Tyr Thr Phe Cys Lys Ala Phe Pro Phe His Ile Ile Phe Asp  
 Arg Asp Leu Val Val Thr Gln Cys Gly Asn Ala Ile Tyr Arg Val Leu  
 Pro Gln Leu Gln Pro Gly Asn Cys Ser Leu Leu Ser Val Phe Ser Leu  
 Val Arg Pro His Ile Asp Ile Ser Phe His Gly Ile Leu Ser His Ile  
 Asn Thr Val Phe Val Leu Arg Ser Lys Glu Gly Leu Leu Asp Val Glu  
 Lys Leu Glu Cys Glu Asp Glu Leu Thr Gly Thr Glu Ile Ser Cys Leu  
 Arg Leu Lys Gly Gln Met Ile Tyr Leu Pro Glu Ala Asp Ser Ile Leu  
 Phe Leu Cys Ser Pro Ser Val Met Asn Leu Asp Asp Leu Thr Arg Arg  
 Gly Leu Tyr Leu Ser Asp Ile Pro Leu His Asp Ala Thr Arg Asp Leu  
 Val Leu Leu Gly Glu Gln Phe Arg Glu Glu Tyr Lys Leu Thr Gln Glu  
 Leu Glu Ile Leu Thr Asp Arg Leu Gln Leu Thr Leu Arg Ala Leu Glu  
 Asp Glu Lys Lys Lys Thr Asp Thr Leu Leu Tyr Ser Val Leu Pro Pro  
 Ser Val Ala Asn Glu Leu Arg His Lys Arg Pro Val Pro Ala Lys Arg  
 Tyr Asp Asn Val Thr Ile Leu Phe Ser Gly Ile Val Gly Phe Asn Ala  
 Phe Cys Ser Lys His Ala Ser Gly Glu Gly Ala Met Lys Ile Val Asn  
 Leu Leu Asn Asp Leu Tyr Thr Arg Phe Asp Thr Leu Thr Asp Ser Arg  
 Lys Asn Pro Phe Val Tyr Lys Val Glu Thr Val Gly Asp Lys Tyr Met  
 Thr Val Ser Gly Leu Pro Glu Pro Cys Ile His His Ala Arg Ser Ile  
 Cys His Leu Ala Leu Asp Met Met Glu Ile Ala Gly Gln Val Gln Val  
 Asp Gly Glu Ser Val Gln Ile Thr Ile Gly Ile His Thr Gly Glu Val  
 Val Thr Gly Val Ile Gly Gln Arg Met Pro Arg Tyr Cys Leu Phe Gly  
 Asn Thr Val Asn Leu Thr Ser Arg Thr Glu Thr Thr Gly Glu Lys Gly  
 Lys Ile Asn Val Ser Glu Tyr Thr Tyr Arg Cys Leu Met Ser Pro Glu  
 Asn Ser Asp Pro Gln Phe His Leu Glu His Arg Gly Pro Val Ser Met  
 Lys Gly Lys Lys Glu Pro Met Gln Val Trp Phe Leu Ser Arg Lys Asn  
 Thr Gly Thr Glu Glu Thr Lys Gln Asp Asp Asp end

**Figure 22**

Phe Thr Pro Arg Ser Arg Glu Glu Leu Pro Pro Asn Phe Pro

**Figure 23**

Lys Gly Lys Lys Glu Pro Met Gln Val Trp Phe Leu Ser Arg Lys Asn  
Thr Gly Thr Glu Glu Thr

**Figure 24**

upper primer

AAAAGGATCC ATGTTCTGCA CGAAGCTC

lower primer

ATTATGGAAG CAGGGAGG


**Figure 25**

upper primer


AAAAGGATCC ATGTACGGAT TTGTGAAT


lower primer


ATGCGTGATT CCTGGGTACC


**Figure 26**



**Figure 27**

**Figure 28**

**Figure 29**

**Figure 30**

<b>DECLARATION FOR UTILITY OR DESIGN PATENT APPLICATION</b> <b>(37 CFR 1.63)</b>	Attorney Docket Number	VOS-101
	First Named Inventor	SCHMIDT, Harald
	<b>COMPLETE IF KNOWN</b>	
	Application Number	09/762,767
	Filing Date	August 16, 1999
	Group Art Unit	
<input type="checkbox"/> Declaration Submitted with Initial Filing <input checked="" type="checkbox"/> Declaration Submitted after Initial Filing	Examiner Name	

As a below named inventor, I hereby declare that:

My residence, post office address, and citizenship are as stated below next to my name.

I believe I am the original, first, and sole inventor (if only one name is listed) or an original, first, and joint inventor (if more than one name is listed) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

the specification of which:

☐ is attached hereto;

☒ was filed on on August 16, 1999 as United States Application Number or PCT International Application Number PCT/DE99/02601 and was amended on September 5, 2000.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the patentability of this application, as defined in 37 CFR 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Number(s)	Country	Foreign Filing Date (MM/DD/YYYY)	Priority Not Claimed	Certified Copy Attached?	
				YES	NO
198 37 015.6	Germany	08/14/98	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

☐ Additional foreign application numbers are listed on a supplemental priority data sheet attached hereto.

I hereby claim the benefits under 35 U.S.C. 119(e) of any United States provisional application listed below.

Application Number(s)	Foreign Filing Date (MM/DD/YYYY)	<input type="checkbox"/> Additional provisional application numbers are listed on a supplemental priority data sheet attached hereto.

# DECLARATION

## - Utility or Design Patent Application

I hereby claim the benefit under 35 U.S.C. 120 of any United States application(s), or 365(c) of any PCT international application designating the United States of America, filed below, and insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of Title 35 U.S.C. 112, I acknowledge the duty to disclose information which is material to patentability, as defined in 37 C.F.R. 1.56, which became available between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. Parent Application or PCT Parent Number	Parent Filing Date (MM/DD/YYYY)	Parent Patent Number (if applicable)

☐ Additional U.S. or PCT international application numbers are listed on a supplemental priority data sheet attached hereto.

As a named inventor, I hereby appoint the following registered practitioner(s) to prosecute this application and to transact all business connected therewith in the Patent and Trademark Office:

☐ Customer Number \_\_\_\_\_ or ☒ Registered Practitioner(s) name/registration numbers listed below

Name	Registration Number	Name	Registration Number
Arne M. Olson	30,203	Michael A. Hiert	29,807
Dolores T. Kenney	31,269	Talivaldis Cepuritis	20,818
Seymour Rothstein	19,369	Daniel J. Deneufbourg	33,675

☒ Additional registered practitioner(s) named on supplemental Registered Practitioner Information sheet attached hereto.

Direct all correspondence to: ☐ Customer Number \_\_\_\_\_ or ☒ Correspondence address below

Name <u>CEPURITIS, Talivaldis</u> <u>OLSON &amp; HIERL, LTD.</u>			
Address <u>20 North Wacker Drive, 36th Floor</u>			
City <u>Chicago</u>	State <u>IL</u>	ZIP <u>60606</u>	
Country <u>US</u>	Telephone <u>(312) 580-1180</u>	Fax <u>(312) 580-1189</u>	

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements are made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Name of Sole or First Inventor:		<input type="checkbox"/> A petition has been filed for this unsigned inventor	
Given Name (first and middle, if any)		Family Name or Surname	
<u>Harald</u>		<u>SCHMIDT</u>	
Inventor's signature	<u>[Signature]</u>		Date: <u>4/3/02</u>
Residence	City <u>Wienburg</u> <u>Gießen</u>	State	Country DE Citizenship DE
Post Office Address	<u>Langer Gäßchen 26</u> <u>Arndtstr. 14</u>		
City <u>Wienburg</u> <u>Gießen</u>	State	<del>ZIP 97078 DE</del> <u>ZIP 35392 DE</u>	Citizenship DE
<input checked="" type="checkbox"/> Additional inventors are being named on the <u>1</u> supplemental Additional Inventor(s) Sheet(s) attached hereto			

## DECLARATION

ADDITIONAL INVENTOR(S)  
Supplemental Sheet

Page 1 of 1

<b>Name of Additional Joint Inventor, if any:</b>		<input type="checkbox"/> A petition has been filed for this unsigned inventor		
Given Name (first and middle, if any)		Family Name or Surname		
2- <u>Ulrike</u>		<u>ZABEL</u>		
Inventor's signature	<u>U. Zabel</u>		Date: <u>4/27/01</u>	
Residence	City <u>Rottendorf</u>	State	Country DE	Citizenship DE <b>DEX</b>
Post Office Address	<u>Am Seelein 26</u>			
	City <u>Rottendorf</u>	State	ZIP <u>97228</u>	Citizenship DE
<b>Name of Additional Joint Inventor, if any:</b>		<input type="checkbox"/> A petition has been filed for this unsigned inventor		
Given Name (first and middle, if any)		Family Name or Surname		
3- <u>Wolfgang</u>		<u>POLLER</u>		
Inventor's signature	<u>W. Poller</u>		Date: <u>5/8/01</u>	
Residence	City <u>Berlin</u>	State	Country DE	Citizenship DE <b>DEX</b>
Post Office Address	<u>Hauptstrasse 101 Rohrbecker Weg 48c</u>			
	City <u>Berlin</u>	State	ZIP <u>12469</u>	Citizenship DE
<b>Name of Additional Joint Inventor, if any:</b>		<input type="checkbox"/> A petition has been filed for this unsigned inventor		
Given Name (first and middle, if any)		Family Name or Surname		
Inventor's signature			Date:	
Residence	City	State	Country	Citizenship
Post Office Address				
	City	State	ZIP	Citizenship

DECLARATION		REGISTERED PRACTITIONER INFORMATION (Supplemental Sheet)	
Name	Reg. No.	Name	Reg. No.
Joseph M. Kuo	<u>38,943</u>		
Martin J. Corn	<u>35,847</u>		
Richard L. Robertson	<u>31,415</u>		
David A. Gottardo	<u>46,736</u>		
Robert J. Ross	<u>45,058</u>		



SEQUENCE LISTING

<110> SCHMIDT, Harald  
ZABEL, Ulrike  
POLLER, Wolfgang

<120> Isolated and Purified Human Soluble  
Guanylylcyclase alpha 1/beta 1 (hsGCalpha/beta 1)

<130> VOS-101

<140> US 09/762,767

<141> 2001-02-13

<150> PCT/DE99/02601

<151> 1999-08-16

<150> DE 198 37 015.6

<151> 1998-08-14

<160> 10

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 3015

<212> DNA

<213> genomic DNA

<400> 1  
cccttatggc gattggcgcg ctgcagagac caggactcag ttccctgcc ctagtctgag 60  
cctagtgggt ggaactcagc tcagagtcag ttttcagaa gcagggttca gtgcagagtt 120  
ttcctacact ttctctcgcg tagagcagcg agcagcctgg aacagaccca ggccggaggac 180  
acctgtgggg gagggagcgc ctggaggagc ttagagaccc cagccggggc gtatctcacc 240  
atgtgcggat ttgcgaggcg cgccttggag ctgctagaga tccggaagca cagccccgag 300  
gtgtgcgaag ccaccaagac tgcggctctt ggagaaagcg tgagcagggg gccaccgcgg 360  
ttctcgcgcc tgctgcacc ctgtgcctcg agctgcctga cagtgaacaat gacatcccag 420  
ttaccagtggt ccttgaattg atagtggctt ctgtttgtca gtctcatata agaactacag 480  
ctcatcagga ggagatcgca gcagggttaag agacaccaac accatgttct gcacgaagct 540  
caaggatctc aagatcacag gagagtgctc ttctctctta ctggcaccag gtccaagtcc 600  
taacagagtt tcagaggagg cagcagggaag ctcaagagag tgcaaaagcaa ccgtgcccatt 660  
ctgtcaagac attcctgaga agaacataca agaaagtctt cctcaaaagaa aaaccagctg 720  
gagccgagtc tatcttcaca ctttggcaga gagtatttgc aaactgattt tcccagagtt 780  
tgaacggctg aatgtttcac ttcagagaaac attggcaag cacaataataa aagaaagcag 840  
gaaatctttg gaagagagaag actttgaaaa aacaattgca gagcaagcag ttgcagcagg 900  
aggtccagtg gaggttatca aagaatctct tggtagagag gtttttaaaa tatgttacga 960  
ggaagatgaa aacatccttg ggggtggttg aggcaccctt aaagattttt taacacagctt 1020  
cagtagccct ctgaaacaga gcagccattg ccaagaagca ggaaaaaggg gcagggttga 1080  
ggacgcctcc attctatgcc tggataagga ggatgattt ctacatgttt actactcttt 1140  
ccctaagaga accacctccc tgattcttcc cggcatcata aaggcagctg ctacagctatt 1200  
atatgaaacg gaagtggaa tgctgttaat gcctccctgc ttccataatg attgcagcga 1260  
gtttgtgaat cagccctact tgttgtactc cgttcacatg aaaagcacca agccatccct 1320

```

gtccccccagc aaaccccagc cctcgctggc gattcccaca tcgctattct gcaagacatt 1380
tccatttccat ttcattgttg acaaagatat gacaattctg caatttggca atggcatcag 1440
aaggctgtagt aacaggagag actttcaagg aaagcctaata tttgaagaat actttgaaat 1500
tctgactcca aaaatcaacc agacgttttag cgggatcatg actatgttga atatgcatgt 1560
tggtgtacga gtgaggagat gggacaactc tgtgaagaaa tcttcaagggt tbatggacct 1620
caaaaggcca atgatctaca ttgttgatc cagtgcatac ttgtttttgg ggctaccctg 1680
tgtggacaga ttagaagatt ttacaggacg agggctctac ctctcagaca tcccaattca 1740
caatgcactg agggatgtgg tcttaatagg ggaacaagcc cgagctcaag atggcctgaa 1800
gaagaggctg gggaaagctga aggcctaccct tgagcaagcc caccagcccc tggaggagga 1860
gaagaaaaag cacagtagacc ttctgtgctc catatttccc tgtgaggttg ctgcagcagt 1920
gtggcaaggg caagtgtgac aagccaagaa gtccagtaat gtcaccaatg ctccttcaaga 1980
catcgctggg ttcactgcca tctgctccca gtgctcaccg ctgcaggtag tcaccatgct 2040
caatgcactg tacactcgct tgcaccagca gtgtggagag ctggatgtct acaagggtga 2100
gaccattggc gatgcctatt gtgtagctgg gggattacac aaagagagtg atactcatgc 2160
tgttccagata gcgctgatgg cctgaagat gatggagctc tctgatgaag ttatgtctcc 2220
ccatggagaa cctatcaaga tgcgaattgg actgcactct ggatcagttt ttgctggcgt 2280
cggtggagtt aaaaatgcccc gttactgtct ttttggaaac aatgtcactc tggctaacca 2340
atttgagctc tgcaggtgac cagcaaaaaa caatgtcagc caaccaactt agcagattact 2400
caaaactgt cctggtttcc tggttaccctc tcgatcaagg gaggaacttc caccaaactt 2460
ccctagtga atccccggaa tctgccattt tctggatgct taccacaacg gaacaaactc 2520
aaaacatgc ttccaaaaga aagatgtgga agatggcaat gccaattttt taggcaaacg 2580
atcagggaata gattagcaac ctataacct attataagt ctttgggggt tgactcattg 2640
aagatgtgta gagcctctga aagcacttta gggattgtag atggctaaca agcagattta 2700
aaatttcagg agccaagtca caatctttct cctgtttaac atgcaaaaat gtactcactt 2760
cagtcattca gctcttcaag aaaaaaaa aaacctttaa aagctacttt tgtggaggta 2820
ttctattat ataaccagca ctactacct gtactcaaaa ttcagcaccct tgatcatata 2880
tcagataaatt gttagcaatt gtacaaactg atggagtcac ctgcaactct atactctggt 2940
ggaatgccaat ggttattaaa gtgtgtttgt gatagtgtcg tcaaaaaaaa aaaaaaaa 3000
aaaaaaaaaaaa 3015

```

&lt;210&gt; 2

&lt;211&gt; 690

&lt;212&gt; PRT

&lt;213&gt; human soluble guanylyl cyclase a1 (hsGCal)

&lt;400&gt; 2

```

Met Phe Cys Thr Lys Leu Lys Asp Leu Lys Ile Thr Gly Glu Cys Pro
1 5 10 15
Phe Ser Leu Leu Ala Pro Gly Gln Val Pro Asn Glu Ser Ser Glu Glu
20 25 30
Ala Ala Gly Ser Ser Glu Ser Cys Lys Ala Thr Val Pro Ile Cys Gln
35 40 45
Asp Ile Pro Glu Lys Asn Ile Gln Glu Ser Leu Pro Gln Arg Lys Thr
50 55 60
Ser Arg Ser Arg Val Tyr Leu His Thr Leu Ala Glu Ser Ile Cys Lys
65 70 75 80
Leu Ile Phe Pro Glu Phe Glu Arg Leu Asn Val Ala Leu Gln Arg Thr
85 90 95
Leu Ala Lys His Lys Ile Lys Glu Ser Arg Lys Ser Leu Glu Arg Glu
100 105 110
Asp Phe Glu Lys Thr Ile Ala Glu Gln Ala Val Ala Ala Gly Val Pro
115 120 125
Val Glu Val Ile Lys Glu Ser Leu Gly Glu Glu Val Phe Lys Ile Cys
130 135 140
Tyr Glu Glu Asp Glu Asn Ile Leu Gly Val Val Gly Gly Thr Leu Lys

```

145					150					155					160
Asp	Phe	Leu	Asn	Ser	Phe	Ser	Thr	Leu	Leu	Lys	Gln	Ser	Ser	His	Cys
				165					170					175	
Gln	Glu	Ala	Gly	Lys	Arg	Gly	Arg	Leu	Glu	Asp	Ala	Ser	Ile	Leu	Cys
			180					185					190		
Leu	Asp	Lys	Glu	Asp	Asp	Phe	Leu	His	Val	Tyr	Tyr	Phe	Phe	Pro	Lys
		195					200					205			
Arg	Thr	Thr	Ser	Leu	Ile	Leu	Pro	Gly	Ile	Ile	Lys	Ala	Ala	Ala	His
		210				215					220				
Val	Leu	Tyr	Glu	Thr	Glu	Val	Glu	Val	Ser	Leu	Met	Pro	Pro	Cys	Phe
225					230					235				240	
His	Asn	Asp	Cys	Ser	Glu	Phe	Val	Asn	Gln	Pro	Tyr	Leu	Leu	Tyr	Ser
				245					250					255	
Val	His	Met	Lys	Ser	Thr	Lys	Pro	Ser	Leu	Ser	Pro	Ser	Lys	Pro	Gln
			260				265						270		
Ser	Ser	Leu	Val	Ile	Pro	Thr	Ser	Leu	Phe	Cys	Lys	Thr	Phe	Pro	Phe
		275					280					285			
His	Phe	Met	Phe	Asp	Lys	Asp	Met	Thr	Ile	Leu	Gln	Phe	Gly	Asn	Gly
		290				295					300				
Ile	Arg	Arg	Leu	Met	Asn	Arg	Arg	Asp	Phe	Gln	Gly	Lys	Pro	Asn	Phe
305					310					315				320	
Glu	Glu	Tyr	Phe	Glu	Ile	Leu	Thr	Pro	Lys	Ile	Asn	Gln	Thr	Phe	Ser
			325						330					335	
Gly	Ile	Met	Thr	Met	Leu	Asn	Met	Gln	Phe	Val	Val	Arg	Val	Arg	Arg
			340					345					350		
Trp	Asp	Asn	Ser	Val	Lys	Lys	Ser	Ser	Arg	Val	Met	Asp	Leu	Lys	Gly
		355					360					365			
Gln	Met	Ile	Tyr	Ile	Val	Glu	Ser	Ser	Ala	Ile	Leu	Phe	Leu	Gly	Ser
		370				375					380				
Pro	Cys	Val	Asp	Arg	Leu	Glu	Asp	Phe	Thr	Gly	Arg	Gly	Ile	Tyr	Leu
385					390					395				400	
Ser	Asp	Ile	Pro	Ile	His	Asn	Ala	Leu	Arg	Asp	Val	Val	Leu	Ile	Gly
				405					410					415	
Glu	Gln	Ala	Arg	Ala	Gln	Asp	Gly	Leu	Lys	Lys	Arg	Leu	Gly	Lys	Leu
			420					425					430		
Lys	Ala	Thr	Leu	Glu	Gln	Ala	His	Gln	Ala	Leu	Glu	Glu	Gly	Lys	Lys
		435				440						445			
Lys	Thr	Val	Asp	Leu	Leu	Cys	Ser	Ile	Phe	Pro	Cys	Glu	Val	Ala	Gln
		450				455					460				
Gln	Leu	Trp	Gln	Gly	Gln	Val	Val	Gln	Ala	Lys	Lys	Phe	Ser	Asn	Val
465					470					475				480	
Thr	Met	Leu	Phe	Ser	Asp	Ile	Val	Gly	Phe	Thr	Ala	Ile	Cys	Ser	Gln
				485					490					495	
Cys	Ser	Pro	Leu	Gln	Val	Ile	Thr	Met	Leu	Asn	Ala	Leu	Tyr	Thr	Arg
			500												

```

Arg Tyr Cys Leu Phe Gly Asn Asn Val Thr Leu Ala Asn Lys Phe Glu
      595                      600                      605
Ser Cys Ser Val Pro Arg Lys Ile Asn Val Ser Pro Thr Thr Tyr Arg
      610                      615                      620
Leu Leu Lys Asp Cys Pro Gly Phe Val Phe Thr Pro Arg Ser Arg Glu
625                      630                      635                      640
Glu Leu Pro Pro Asn Phe Pro Ser Glu Ile Pro Gly Ile Cys His Phe
      645                      650                      655
Leu Asp Ala Tyr Gln Gln Gly Thr Asn Ser Lys Pro Cys Phe Gln Lys
      660                      665                      670
Lys Asp Val Glu Asp Gly Asn Ala Asn Phe Leu Gly Lys Ala Ser Gly
      675                      680                      685
Ile Asp
      690

```

```

<210> 3
<211> 2443
<212> DNA
<213> genomic DNA

```

```

<400> 3
cccccccgcg cgcgtgccgc ctctgcctgg gtcctctcgg ccgtacctct gcgtgggggc 60
tgctcccccgc gctcccgggt cagacacccat gtacgggattt gtgaatcacg ccttgaggatt 120
gctgggtgatc gcgaattacg gccccgagggt gtgggaagac atcaaaaaag aggcacaggt 180
agattgaagaa ggacagtttc ttgtcagaat aatatatgat gactccaaaaa cttatgattt 240
ggttgctgctc gcaagcaaaag tctcaaatc caatgctgga gaaatcctcc aaatgtttgg 300
gaagatgtttt tctgcttttt gccaaagaatc tggttatgat acaatcttgc gtgtcctggg 360
ctctaagtgc agagaatttc tacagaacct tgaatgctcg cagcacccac ttgctaccat 420
ctacccaggaa atgcgtgcac ctctccttag gtgcactgat gcagaaaaag gcaaaggact 480
cattttgcac tactactcag agagagaagg acttcaggat attgtcattg gaatcatcaa 540
aacagtggca caacaaatcc atggcactga aatagacatg aagggtattc agcaaaagaa 600
tgaagaatgt gatcatactc aatttttaat tgaagaaaaa gagtcaaaaag aagaggattt 660
ttatgaagat cttgacagat ttgaagaaaaa tggtagccag gaatcacgca tcagcccata 720
tacattctgc aaagcttttc cttttcataa aatatttgac cgggacctag tggctactca 780
gtgtggcaat gctatataca gagttctccc ccagctccag cctgggaatt gcagccttct 840
gtctgtcttc tctcgggttc gtctctcatat tgatattagt ttccatggga tcttttctca 900
catcaatact gtttttgat tgaagaagcaa ggaaggattg ttggatgtgg agaaaattaga 960
atgtgaggat gaactgactg ggaactgagat cagctgctta cgtctcaagg gtcaaatgat 1020
ctacttacct gaagcagata gcatactttt tctatgttca ccaagtgtca tgaacctgga 1080
cgatttgaca aggaagagggc tgtatctaa gtagactcct ctgcatgatg ccacgcgcga 1140
tcttctcttc ttggggagaa aatttagaga ggaatacaaa ctccaccaag aactggaaat 1200
cctcaactgc aggtctacagc tcacgttaag agccctggaa gatgaaaaga aagaacacaga 1260
cacattgctg tattctgtcc ttccctcgcg tgttgccaat gagctgcgag acaagcgtcc 1320
agtgctctgcc aaaagatatg acaatgtgac catcctcttt agtggcattg tgggcttcaa 1380
tgctttctgt agcaagcatg catctggaga aggagccatg aagatcgta accctctcaa 1440
cgacctctac accagatttg acacactgac tgattccccg aaaaacccat ttggttataa 1500
gggtggagact gttggtgaca agtatatgac agtgagtggt ttaccagagc catgacttca 1560
ccatgcacga tccatctgcc acctggcctt ggacatgatg gaaattgctg gccaggttca 1620
agtagatggt gaactctgtc agataacaat agggatatac actggagaggt tagttacagg 1680
tgtcatagat cagcgggatgc ctcgatactg tctttttggg aatactgtca acctcacaag 1740
ccgaacagaa accacaggag aaaaagggaa aataaatatg tctgaatata acctcagagt 1800
tcttatgtct ccagaaaatt cagatccaca attccacttg gagcacagag gccccagtgc 1860
catgaagggc aaaaaagaac caatgcaagt ttggtttcta tccagaaaaa atacaggaac 1920
agaggaaca aagcaggatg atgactgaat ctggattat ggggtgaaga ggagtacaga 1980

```

```

ctaggttcca gttttctcct aacacgtgcc aagcccagga gcagttcttc cctatggata 2040
cagattttct tttgtccttg tccattaccc caagactttc ttctagatat atctctcact 2100
atccgttatt caaccttagc tctgctttct attacttttt aggtcttagt atattatcta 2160
aagtttggct tttgatgtgg atgatgtgag cttcatgtgt cttaaaatct actacaagca 2220
ttacctaaaca tgggtgatctg caagtagtag gcaccaata aatatttgggt gaatttagtt 2280
aaatgaaact gaacagtggt tggccatgtg tatatttata tcatgtttac caaatctggt 2340
tagtgttcca catatatgta tatgtatatt ttaatgacta taatgtaata aagtttatat 2400
catgttggtg tatatcatta tagaaatcat tttctaaagg agt 2443

```

&lt;210&gt; 4

&lt;211&gt; 619

&lt;212&gt; PRT

&lt;213&gt; (human soluble guanylyl cyclase b1 (hsGCB1))

&lt;400&gt; 4

```

Met Tyr Gly Phe Val Asn His Ala Leu Glu Leu Leu Val Ile Arg Asn
  1          5          10          15
Tyr Gly Pro Glu Val Trp Glu Asp Ile Lys Lys Glu Ala Gln Leu Asp
  20          25          30
Glu Glu Gly Gln Phe Leu Val Arg Ile Ile Tyr Asp Asp Ser Lys Thr
  35          40          45
Tyr Asp Leu Val Ala Ala Ala Ser Lys Val Leu Asn Leu Asn Ala Gly
  50          55          60
Glu Ile Leu Gln Met Phe Gly Lys Met Phe Phe Val Phe Cys Gln Glu
  65          70          75          80
Ser Gly Tyr Asp Thr Ile Leu Arg Val Leu Gly Ser Asn Val Arg Glu
  85          90          95
Phe Leu Gln Asn Leu Asp Ala Leu His Asp His Leu Ala Thr Ile Tyr
  100          105          110
Pro Gly Met Arg Ala Pro Ser Phe Arg Cys Thr Asp Ala Glu Lys Gly
  115          120          125
Lys Gly Leu Ile Leu His Tyr Tyr Ser Glu Arg Glu Gly Leu Gln Asp
  130          135          140
Ile Val Ile Gly Ile Ile Lys Thr Val Ala Gln Gln Ile His Gly Thr
  145          150          155          160
Glu Ile Asp Met Lys Val Ile Gln Gln Arg Asn Glu Glu Cys Asp His
  165          170          175
Thr Gln Phe Leu Ile Glu Glu Lys Glu Ser Lys Glu Glu Asp Phe Tyr
  180          185          190
Glu Asp Leu Asp Arg Phe Glu Glu Asn Gly Thr Gln Glu Ser Arg Ile
  195          200          205
Ser Pro Tyr Thr Phe Cys Lys Ala Phe Pro Phe His Ile Ile Phe Asp
  210          215          220
Arg Asp Leu Val Val Thr Gln Cys Gly Asn Ala Ile Tyr Arg Val Leu
  225          230          235          240
Pro Gln Leu Gln Pro Gly Asn Cys Ser Leu Leu Ser Val Phe Ser Leu
  245          250          255
Val Arg Pro His Ile Asp Ile Ser Phe His Gly Ile Leu Ser His Ile
  260          265          270
Asn Thr Val Phe Val Leu Arg Ser Lys Glu Gly Leu Leu Asp Val Glu
  275          280          285
Lys Leu Glu Cys Glu Asp Glu Leu Thr Gly Thr Glu Ile Ser Cys Leu
  290          295          300
Arg Leu Lys Gly Gln Met Ile Tyr Leu Pro Glu Ala Asp Ser Ile Leu
  305          310          315          320

```

Phe Leu Cys Ser Pro Ser Val Met Asn Leu Asp Asp Leu Thr Arg Arg  
 325 330 335  
 Gly Leu Tyr Leu Ser Asp Ile Pro Leu His Asp Ala Thr Arg Asp Leu  
 340 345 350  
 Val Leu Leu Gly Glu Gln Phe Arg Glu Glu Tyr Lys Leu Thr Gln Glu  
 355 360 365  
 Leu Glu Ile Leu Thr Asp Arg Leu Gln Leu Thr Leu Arg Ala Leu Glu  
 370 375 380  
 Asp Glu Lys Lys Lys Thr Asp Thr Leu Leu Tyr Ser Val Leu Pro Pro  
 385 390 395 400  
 Ser Val Ala Asn Glu Leu Arg His Lys Arg Pro Val Pro Ala Lys Arg  
 405 410 415  
 Tyr Asp Asn Val Thr Ile Leu Phe Ser Gly Ile Val Gly Phe Asn Ala  
 420 425 430  
 Phe Cys Ser Lys His Ala Ser Gly Glu Gly Ala Met Lys Ile Val Asn  
 435 440 445  
 Leu Leu Asn Asp Leu Tyr Thr Arg Phe Asp Thr Leu Thr Asp Ser Arg  
 450 455 460  
 Lys Asn Pro Phe Val Tyr Lys Val Glu Thr Val Gly Asp Lys Tyr Met  
 465 470 475 480  
 Thr Val Ser Gly Leu Pro Glu Pro Cys Ile His His Ala Arg Ser Ile  
 485 490 495  
 Cys His Leu Ala Leu Asp Met Met Glu Ile Ala Gly Gln Val Gln Val  
 500 505 510  
 Asp Gly Glu Ser Val Gln Ile Thr Ile Gly Ile His Thr Gly Glu Val  
 515 520 525  
 Val Thr Gly Val Ile Gly Gln Arg Met Pro Arg Tyr Cys Leu Phe Gly  
 530 535 540  
 Asn Thr Val Asn Leu Thr Ser Arg Thr Glu Thr Thr Gly Glu Lys Gly  
 545 550 555 560  
 Lys Ile Asn Val Ser Glu Tyr Thr Tyr Arg Cys Leu Met Ser Pro Glu  
 565 570 575  
 Asn Ser Asp Pro Gln Phe His Leu Glu His Arg Gly Pro Val Ser Met  
 580 585 590  
 Lys Gly Lys Lys Glu Pro Met Gln Val Trp Phe Leu Ser Arg Lys Asn  
 595 600 605  
 Thr Gly Thr Glu Glu Thr Lys Gln Asp Asp Asp  
 610 615

<210> 5  
 <211> 14  
 <212> PRT  
 <213> peptide (amino acids 634-647 of hsgCal)

<400> 5  
 Phe Thr Pro Arg Ser Arg Glu Glu Leu Pro Pro Asn Phe Pro  
 1 5 10

<210> 6  
 <211> 22  
 <212> PRT  
 <213> peptide (amino acids 593-614 of hsgCbl)

&lt;400&gt; 6

Lys Gly Lys Lys Glu Pro Met Gln Val Trp Phe Leu Ser Arg Lys Asn

1

5

10

15

Thr Gly Thr Glu Glu Thr

20

&lt;210&gt; 7

&lt;211&gt; 28

&lt;212&gt; DNA

&lt;213&gt; genomic DNA

&lt;400&gt; 7

aaaaggatcc atgttctgca cgaagctc

28

&lt;210&gt; 8

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; genomic DNA

&lt;400&gt; 8

attatggaag cagggagg

18

&lt;210&gt; 9

&lt;211&gt; 28

&lt;212&gt; DNA

&lt;213&gt; genomic DNA

&lt;400&gt; 9

aaaaggatcc atgtacggat ttgtgaat

28

&lt;210&gt; 10

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; genomic DNA

&lt;400&gt; 10

atgcgtgatt cctgggtacc

20

09/762767

PTO/PCT Rec'd 01 JUN 2001

#3

1

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANTS:

SCHMIDT, Harald

ZABEL, Ulrike

POLLER, Wolfgang

(ii) TITLE OF THE INVENTION: Isolated and purified human soluble guanylyl cyclase  $\alpha 1/\beta 1$  (hsGCa1/ $\beta 1$ )

(iii) VOS-101

(iv) US 09/762,767  
2001-02-13(iv) PCT/DE99/02601  
1999-08-16(iv) DE 198 37 015.6  
1998-08-14

(v) NUMBER OF SEQUENCES: 10

## (vi) COMPUTER-READABLE FORM:

(A) MEDIUM TYPE: Floppy disk

(B) COMPUTER: IBM PC compatible

(C) OPERATING SYSTEM: PC-DOS/MS-DOS

(D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPA)

## (2) DATA TO SEQ ID NO: 1:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 3015 basepairs

(B) TYPE: nucleotide

(C) STRANDEDNESS: doublestrand

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: genomic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

CCCTTATGGC	GATTGGGCGG	CTGCAGAGAC	CAGGACTCAG	TTCCCTTGCC	CTAGTCTGAG	60
CCCTAGTGGG	GGGACTCAGC	TCAGAGTCAG	TTTCCAGAA	GCAGGTTTCA	GTGCAGAGTT	120
TTCTCTACCT	TTTCTCGCGC	TAGAGCAGCG	AGCAGCCTGG	AACAGACCCA	GGCCGAGGAC	180
ACCTGTGGGG	GAGGGAGCGC	CTGGAGGAGC	TTAGAGACCC	CAGCCGGGCG	TGATCTCACC	240
ATGTGCGGAT	TTGCGAGGCG	CGCCCTGGAG	CTGCTAGAGA	TCCGGAAGCA	CAGCCCCGAG	300
GTGTGCGAAG	CCACCAAGAC	TGCGGCTCTT	GGAGAAAGCG	TGAGCAGGGG	GCACCCGCGG	360
TCTCCGCGCC	TGCTCGCACC	CTGTGCGCTG	AGCTGCCTGA	CAGTGACAAT	GACATCCGAG	420
TTACCACTGT	CCTTGAATTG	ATAGTGCGCT	CTGTTTGTC	GTCTCATATA	AGAACTACAG	480
CTCATCAGGA	GGAGATCGCA	GCAGGGTAAG	AGACCCAAC	ACCATGTTCT	GCACCAAGCT	540
CAAGGATCTC	AAGATCAGAG	GAGAGTGTCC	TTTCTCCTTA	CTGGCACCAG	GTCAGGTTCC	600
CTCATCAGGA	TCAGAGGAGG	CAGCAGGAAG	CTCAGAGAGC	TGCAAGCAAA	CCGTGCCCAT	660
CTGTCAAGAC	ATTCTTGAGA	AGACATACAA	AGAAAGTCTT	CCTCAAGAAA	AAACCAAGTC	720
GAGCCGAGTC	TATCTTACAA	CTTTGGCAGA	GAGTATTTGC	AAACTGATT	TCCGAGAGTT	780
TGAACGGGTC	AATGTTGCAC	TTACAGAAAC	ATTGGCAAG	CACAAAATAA	AAGAAGAGCA	840
GGAATCTTTG	GAAGAGAAG	ACTTTGAAAA	AACRAATTGC	GAGCAAGCAG	TTGCAGCAGG	900
AGTTCAGGTC	GAGGTTATCA	AGAAATCTCT	TGGTGAAGAG	GTTTTTAAAA	TATGTTACGA	960
GGAAAGATGA	AACATCCTTG	GGGTGGTTGG	AGGCACCCCT	AAAGATTTTT	TAAACAGCTT	1020
CAGTACCCCT	CTGAAACAGA	GCAGCCATTG	CCAGAAAGCA	GGAAAAAGGG	GCAGGCTTGA	1080
GGAGCCCTCC	ATTCATAGGC	TGGATAAGGA	GGATGATTTT	CTACATGTTT	ACTACTTCTT	1140
CCCTAAGAGA	ACCACCTTCC	TGATTTCTCC	CGGCATCATA	TAGGACGCTG	CTCAGCTATT	1200
ATATGAAGAC	GAAAGTGAAG	TGTCGTTAAT	GCCTCCCTGC	TGCATAATG	ATTGGACGGA	1260
GTTTGTGAAT	CAGCCCTACT	TGTTGTACTC	CGTTCACATG	AAAAGCACCA	AGCCATCCCT	1320
GTCCCCCAGC	AAACCCAGT	CCTCGTGGT	GATTCACACA	TGCTATTCT	GCAAGACATT	1380



	TCCATTCAT	TTCAIGTTTG	ACAAAGATAT	GAAATTCCTG	CAATTGTGGCA	ATGGCATCTAG	1440
	AAGGCTGTAG	AACAGGAGAG	ACTTTTCAGG	AAAGGCTAAT	TTTGAAGAAAT	ATCTTTGAAT	1500
	TCTGACTCCA	AAATCAACCC	AGACTTCATG	CGGGATCATG	ACTATGTTGA	ACTTGACAGTT	1560
	TGTTGTGCTGA	GTGAGGAGAT	GGGACCACTC	TGTGAAGAAA	TCITTTCAAGGG	TATGAGCACT	1620
5	CAAGAGGCCAA	ATGATCTACA	TGTTGTAATC	CAGTGCAAAT	TGTTTATTTGG	GGTCAACCCGT	1680
	TGTGACAGAGA	TTAGAAGATT	TTCACGAGAC	AGGGCTCTAC	CTCTCAGACA	TCCCAATCAA	1740
	CAATGCATCT	AGGGATGTGG	TCITTAATAGG	GGAACACGCT	CGAGCTCAAG	ATGCGCTGAA	1800
	GAGAGGGCTG	GGGAGCTGA	AGGCTTACCTT	TGAGCAAGCC	CACCAAGGCC	TGGAGGAGGA	1860
	GAGAAAAAG	ACAGTAGACC	TTCTGTGCTC	CATATTTCCC	TGTGAGGTTG	CTCAGCAGCT	1920
10	GTGGCAGAGG	CAAGTGTGTC	AAGCACAAGAA	TGTCAGTAAT	GTCACCATGC	TCTTCTCAGA	1980
	CATCGTGTGG	TTCACTGCTGA	TCTGCTCCCCA	GTCGTCAACG	CTCAGGGTGA	TCCACCATGTT	2040
	CATAGCATCT	TACACTCGCT	TCGACCAACG	GTGTGGAGAG	CTGGATGTCT	ACAGAGTGGG	2100
	GACCATTTGGC	GATGCTTAAT	GCTGTAGCTG	GGGATATCAC	AAGAGAGTGT	ATATCATCTC	2160
	TGTTTCAGATA	CGCGTGAATG	CGCTGAAGAT	GATGGAGCTC	TCGTGATGA	TATGTCTGCT	2220
15	CCATGGAGAA	CCTATCAAGA	TGCGAATTTG	ACTGCATCT	GAATCAGTTT	TGTCTGGCGT	2280
	CGTTGGAGTT	TAAGATGCCCC	GTTACTGTCT	TTTGTGAAAC	AGATCTCACT	TGGCTAACAA	2340
	ATTTGAGTTC	TGCTGTGTAC	CGACAAAAAT	CAATGTGACG	CCAAACAAC	ACAGATTACT	2400
	CAAAAGCTAT	CCTGGTTTGG	TGTTTACCCC	TCGATCAAGG	GAGGAACCTC	GACCAAACTT	2460
20	CCCTAGTAGT	ATCCCGCCAG	TCTGCCATTT	TCTGGATGCT	TACCAACAAG	GACCAAACTC	2520
	AAACCACTATC	TTCCAAAGAA	AAGATGTGGA	AGATGGCAAT	GCAATTTTTT	TGGGCAAAAG	2580
	ATCAGGAATA	GATTAGCAAC	CTATATACCT	ATTTATAAGT	TTCTGGGGTT	TGACTCACTG	2640
	AAGATGTGTA	GAGCCTCTGA	AAGCACTTTA	GGATTTGTAG	ATGGCTTAACA	AGCAGTATTA	2700
	AAATTTTCAGG	AGCCACTCTCA	CAATCTTTCT	CTCTGTTTAA	ATGACAAAA	GTACTCATTT	2760
	CAGTACTTACG	GCTATCTCAAG	AAAAAAAAAA	AAACCTTAAA	AGCCTACTTT	TGTGGGAGTA	2820
25	TTTCTATTAT	ATAACCAAGTA	CTTACTACTT	TGTACTCAAAA	TTCAGACACT	TGTACATATA	2880
	TCAGATAAAT	TGATGACATC	GTACAAACCT	ATGGAGTCA	CTGCACATCT	ATATCTTGTG	2940
	GGAAATGCCAT	GGTTATTAA	GTGTGTTTGT	GATAGTGTGC	TCAAAAAAAA	AAAAAAAAAA	3000
	AAAAAAAAAA	AAAAA					3015

(2) DATA TO SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 695 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein (human soluble guanylyl cyclase  $\alpha 1$  (hsGC $\alpha 1$ ))

(xi) SEQUENCEDESCRIPTION: SEQ ID NO: 2:

45	Met	Phe	Cys	Thr	Lys <sub>5</sub>	Leu	Lys	Asp	Leu	Lys <sub>10</sub>	Ile	Thr	Gly	Glu	Cys	Pro
	Phe	Ser	Leu	Leu	Ala	Pro	Gly	Gln	Val	Pro	Asn	Glu	Ser	Ser	Glu	Glu
	Ala	Ala	Gly	Ser	Ser	Glu	Ser	Cys	Lys <sub>25</sub>	Ala	Thr	Val	Pro	Ile	Cys	Gln
50	Asp	Ile	Pro	Glu	Lys	Asn	Ile	Gln	Glu	Ser	Leu	Pro	Gln	Arg	Lys	Thr
	Ser	Arg	Ser	Arg	Val	Tyr	Leu	His	Thr	Leu	Ala	Glu	Ser	Ile	Cys	Lys <sub>55</sub>
	Leu	Ile	Phe	Pro	Glu	Phe	Glu	Arg	Leu	Asn	Val	Ala	Leu	Gln	Arg	Thr
55	Leu	Ala	Lys	His	Lys <sub>85</sub>	Ile	Lys	Glu	Ser	Arg	Lys	Ser	Leu	Glu	Arg	Glu
	Asp	Phe	Glu	Lys	Thr	Ile	Ala	Glu	Gln	Ala	Val	Ala	Ala	Gly	Val	Pro
	Val	Glu	Val	Ile	Lys	Glu	Ser	Leu	Gly	Glu	Glu	Val	Phe	Lys	Ile	Cys
60	Val	Glu	Val	Ile	Lys	Glu	Ser	Leu	Gly	Glu	Glu	Val	Phe	Lys	Ile	Cys

Tyr Glu Glu Asp Glu Asn Ile Leu Gly Val Val Gly Gly Thr Leu Lys  
 145 150 155 160  
 Asp Phe Leu Asn Ser Phe Ser Thr Leu Leu Lys Gln Ser Ser His Cys  
 165 170 175  
 5 Gln Glu Ala Gly Lys Arg Gly Arg Leu Glu Asp Ala Ser Ile Leu Cys  
 180 185 190  
 Leu Asp Lys Glu Asp Asp Phe Leu His Val Tyr Tyr Phe Phe Pro Lys  
 195 200 205  
 10 Arg Thr Thr Ser Leu Ile Leu Pro Gly Ile Ile Lys Ala Ala Ala His  
 210 215 220  
 Val Leu Tyr Glu Thr Glu Val Glu Val Ser Leu Met Pro Pro Cys Phe  
 2225 230 235 240  
 His Asn Asp Cys Ser Glu Phe Val Asn Gln Pro Tyr Leu Leu Tyr Ser  
 245 250 255  
 15 Val His Met Lys Ser Thr Lys Pro Ser Leu Ser Pro Ser Lys Pro Gln  
 260 265 270  
 Ser Ser Leu Val Ile Pro Thr Ser Leu Phe Cys Lys Thr Phe Pro Phe  
 275 280 285  
 20 His Phe Met Phe Asp Lys Asp Met Thr Ile Leu Gln Phe Gly Asn Gly  
 290 300 305  
 Ile Arg Arg Leu Met Asn Arg Arg Asp Phe Gln Gly Lys Pro Asn Phe  
 310 315 320 325  
 Glu Glu Tyr Phe Glu Ile Leu Thr Pro Lys Ile Asn Gln Thr Phe Ser  
 330 335 340  
 25 Gly Ile Met Thr Met Leu Asn Met Gln Phe Val Val Arg Val Arg Arg  
 345 350 355  
 Trp Asp Asn Ser Val Lys Lys Ser Ser Arg Val Met Asp Leu Lys Gly  
 360 365 370  
 30 Gln Met Ile Tyr Ile Val Glu Ser Ser Ala Ile Leu Phe Leu Gly Ser  
 375 380 385  
 Pro Cys Val Asp Arg Leu Glu Asp Phe Thr Gly Arg Gly Leu Tyr Leu  
 390 395 400 405  
 Ser Asp Ile Pro Ile His Asn Ala Leu Arg Asp Val Val Leu Ile Gly  
 410 415 420  
 35 Glu Gln Ala Arg Ala Gln Asp Gly Leu Lys Lys Arg Leu Gly Lys Leu  
 425 430 435  
 Lys Ala Thr Leu Glu Gln Ala His Gln Ala Leu Glu Glu Glu Lys Lys  
 440 445 450  
 40 Lys Thr Val Asp Leu Leu Cys Ser Ile Phe Pro Cys Glu Val Ala Gln  
 455 460 465  
 Gln Leu Trp Gln Gly Gln Val Val Gln Ala Lys Lys Phe Ser Asn Val  
 470 475 480 485  
 Thr Met Leu Phe Ser Asp Ile Val Gly Phe Thr Ala Ile Cys Ser Gln  
 490 495 500  
 45 Cys Ser Pro Leu Gln Val Ile Thr Met Leu Asn Ala Leu Tyr Thr Arg  
 505 510 515  
 Phe Asp Gln Gln Cys Gly Glu Leu Asp Val Tyr Lys Val Glu Thr Ile  
 520 525 530  
 50 Gly Asp Ala Tyr Cys Val Ala Gly Gly Leu His Lys Glu Ser Asp Thr  
 535 540 545  
 His Ala Val Gln Ile Ala Leu Met Ala Leu Lys Met Met Glu Leu Ser  
 550 555 560 565  
 Asp Glu Val Met Ser Pro His Gly Glu Pro Ile Lys Met Arg Ile Gly  
 570 575 580  
 55 Leu His Ser Gly Ser Val Phe Ala Gly Val Val Gly Val Lys Met Pro  
 585 590 595  
 Arg Tyr Cys Leu Phe Gly Asn Asn Val Thr Leu Ala Asn Lys Phe Glu  
 600 605 610  
 60 Ser Cys Ser Val Pro Arg Lys Ile Asn Val Ser Pro Thr Thr Tyr Arg  
 615 620 625  
 Leu Leu Lys Asp Cys Pro Gly Phe Val Phe Thr Pro Arg Ser Arg Glu

(2) DATA TO SEO ID NO: 3:

(A) LENGTH: 2443 basepairs

(B) TYPE: nucleotide

(C) STRANDEDNESS: doublestrand

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: genomic DNA

(xi) SEQUENCE DESCRIPTION: SEO ID NO: 3:

CCCCCCCCCG	CGCTGCGCG	CTCTGCCCTGG	GTCCTTCCGG	CGTACCTCT	GGTGGGGGG	60
TGCCCTCCGCG	GCTCCCGGTG	CAGACACCAT	GTACGGATTG	GTGAATCACC	CGCTGGAGTT	120
CTGTGGTTCG	CGCAATTACG	CGCCCGAGT	GTGGGAAGAT	ATCAAAAAAG	ACCGACAGTT	180
AGATGACGAA	GGACAGTTTC	TGTGCAGAT	AATATATGAT	GACTCCAAAA	CTTATGATTT	240
GGTGTCTGCT	CAGACGAAAG	TCTCAATCT	CATCTCGTGA	GAAATCTCTC	GAAGTTTGTG	300
GAAATGATTT	TTCTGCTTTT	CACGAGAATC	TGTTATGATG	ACAACTTTTG	GTGCTCGGG	360
CTCTAATATG	AGAGAAATTC	TACGAAATCT	TGATGCTCTG	CAGCACCACT	TTCGTACCAT	420
CTATCCGAGG	ATGCTGTCAC	CTCTCTTTAG	TGTCACATG	CAGAAAAAGG	CGCAAGGACT	480
TATTTGGCAC	TATGCTCAG	AGAGAGAAAG	ACTTCAGAT	ATTGTCTATT	GAACTCATAA	540
AACAGTGGCA	CAACAAATCC	ATGGCACTGA	AATAGACATG	AAGGTTATTCT	AGCAAGAGAA	600
TGAAGAATGT	GATCATACTG	AATTTTAAAT	TGAAGAAAAA	GAGTCAAAAG	AAGAGGATTT	660
TTATAGAAGT	CTTGACAGAT	TGTAGAAAAT	TGATACCCAG	GAATACGCA	TGCAGCCATA	720
TACATCTCG	AAAGCTTTTC	TTCTTCATAT	AATATTTCAG	CGGGACCTAG	TGGTCACTCA	780
GTGTGGCAAT	GCTATATACA	GAGTTCCTCC	CGAGCTCGAG	CTCGGAATG	CAGCGCTTCT	840
GTCTGCTCTC	TGCGTGGTTC	GTCTCATATG	TGATATTAGT	TTGCATGGGA	TGCTTTCTCA	900
CATCAATACT	GTTTTGTGAT	TGCAAGCAAT	GGAGGAGTAT	TTCGATGTGG	AGCAAAATGA	960
ATGTGAGAGT	GAACTGACGT	GGACTGAGAT	CAGCTGCTTA	CGTCTCAAGG	GTCAAAATGT	1020
CTACTTTACC	GAAGCAGATA	GCATACTTTT	TCTATTTCTT	CCAAGTGTC	TGAACCTGAT	1080
CGATTTGACA	AGGAGAGGGC	TGATATTAAG	TGACATCCCT	CTGCATGATG	CACGCGGCG	1140
TTCTGTCTCT	TTGGGAGAAC	AATTTAGAGA	GGAAATACAA	GTCAACCAAG	TAGTGGAAT	1200
CTCTCATGCG	AGGCTCAGCG	CTCTAGTAG	AGCCCTGGAA	AATGAAAAAG	AAAGTACAGA	1260
CACATGTGCT	TAAATCTGTC	TTCTCCGCTG	TGTTGCCAAT	GAGCTGCGGC	CACAGCGTCC	1320
AGGCGCTGCC	AAAAGATATG	ACAATGTGAC	CATCTCTT	GTGGCGATG	TGCGGTTCAA	1380
TGCTTCTCTG	AGCAAGCATG	CATCTGGAGA	AGGAGCATCA	AGATCTGTCA	ACCTCTCTCA	1440
CGACCTCTAC	ACCAAGATTG	ACACACTGAC	TGATTCCCGG	AAAAACCAT	TTTTTTATA	1500
GGTGGAGACT	GTGTTGTGAC	AGTATATGAC	AGTGAGTGGT	TTACACAGGC	CATGCATCTA	1560
CCATGACAGA	TCCATCTGCC	ACCTGGCCCT	GGACATCTG	GAAATTGCTG	CGCGAGTTCA	1620
AGTAGATAGT	GAACTCTGTC	AGATTAACAT	AGGGATACAC	ACTCGGAGG	TAGTTACAGG	1680
TGTCTAGGA	CGCGGAGTGC	CTCGATACAT	TCITTTTGGG	ATACCTGTCA	ACCTCAAG	1740
CCGACAGCAA	ACCCAGGAGG	AAAGAGGAAA	AATAAATGTG	TCTGAATATA	CATACAGATG	1800
CTTTAATGCT	CAAAAAATCT	CAGTCCACCA	ATTCCATTGT	GAGCAGAAAG	GCCACGTTCT	1860
TCTAAGGTCT	AAAAAAGAAC	CAATCGAAT	TTGGTTCTTG	TCCAGAAAAA	ATCAAGCAAG	1920
AGAGGAAACA	AAGCAGGATG	ACTCTGAAT	CTTGATTATT	GGGGTGAAGA	GAGGTACAGA	1980
CTAGGTTCCA	GTTTTCTCCT	AACACGTGCC	AAGCCGAGGA	CGAGTTCTCT	CCATGTGATTA	2040
CAGATTGTTT	TTTGTCCCTG	TCGATATGCC	CAGACATTTT	CTCTAGATAT	ATTCTCTCAT	2100
ATCCGTTATT	CACCTTAGC	TCTGCTTTCT	ATTACTTTT	AGGCTTATGT	ATATATCTA	2160
AAGTTTGGCT	TTTGTATGTG	ATGATGTGAC	CTTCATGTTG	CTTAAATAGT	ACTACAGCTA	2220

TTACCTAACA	TGGTGATCTG	CAAGTAGTAG	GCACCCAATA	AATATTGTGT	GAATTTAGTT	2280
AAATGAAACT	GAACAGTGTT	TGGCCATGTG	TATATTATTA	TCATGTTTAC	CAAACTCTGT	2340
TAGTGTTTCC	CATATATATGA	TATGATATATT	TTAATGACTA	TAATGATAAG	AAGTTTATAT	2400
CATGTGTGTG	TATATATATGA	TAGAAATCACT	TTTCTAAAGG	AGT		2443

(2) DATA TO SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 619 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein (human soluble guanylyl cyclase b1 (hsGCb1))

(xi) SEQUENCEDESCRIPTION: SEQ ID NO: 4:

20	Met	Tyr	Gly	Phe	Val	Asn	His	Ala	Leu	Glu	Leu	Leu	Val	Ile	Arg	Asn
					5					10					15	
	Tyr	Gly	Pro	Glu	Val	Trp	Glu	Asp	Ile	Lys	Lys	Glu	Ala	Gln	Leu	Asp
				20					25					30		
25	Glu	Glu	Gly	Gln	Phe	Leu	Val	Arg	Ile	Ile	Tyr	Asp	Asp	Ser	Lys	Thr
			35					40					45			
	Tyr	Asp	Leu	Val	Ala	Ala	Ala	Ser	Lys	Val	Leu	Asn	Leu	Asn	Ala	Gly
			50					55					60			
	Glu	Ile	Leu	Gln	Met	Phe	Gly	Lys	Met	Phe	Phe	Val	Phe	Cys	Gln	Glu
			65				70					75			80	
30	Ser	Gly	Tyr	Asp	Thr	Ile	Leu	Arg	Val	Leu	Gly	Ser	Asn	Val	Arg	Glu
				85						90					95	
	Phe	Leu	Gln	Asn	Leu	Asp	Ala	Leu	His	Asp	His	Leu	Ala	Thr	Ile	Tyr
				100					105					110		
35	Pro	Gly	Met	Arg	Ala	Pro	Ser	Phe	Arg	Cys	Thr	Asp	Ala	Glu	Lys	Gly
			115					120					125			
	Lys	Gly	Leu	Ile	Leu	His	Tyr	Tyr	Ser	Glu	Arg	Glu	Gly	Leu	Gln	Asp
			130				135					140				
	Ile	Val	Ile	Gly	Ile	Ile	Lys	Thr	Val	Ala	Gln	Gln	Ile	His	Gly	Thr
			145				150					155			160	
40	Glu	Ile	Asp	Met	Lys	Val	Ile	Gln	Gln	Arg	Asn	Glu	Glu	Cys	Asp	His
				165						170					175	
	Thr	Gln	Phe	Leu	Ile	Glu	Glu	Lys	Glu	Ser	Lys	Glu	Glu	Asp	Phe	Tyr
			180					185					190			
45	Glu	Asp	Leu	Asp	Arg	Phe	Glu	Glu	Asn	Gly	Thr	Gln	Glu	Ser	Arg	Ile
			195					200					205			
	Ser	Pro	Tyr	Thr	Phe	Cys	Lys	Ala	Phe	Pro	Phe	His	Ile	Ile	Phe	Asp
			210				215					220				
	Arg	Asp	Leu	Val	Val	Thr	Gln	Cys	Gly	Asn	Ala	Ile	Tyr	Arg	Val	Leu
			225				230				235				240	
50	Pro	Gln	Leu	Gln	Pro	Gly	Asn	Cys	Ser	Leu	Leu	Ser	Val	Phe	Ser	Leu
				245						250					255	
	Val	Arg	Pro	His	Ile	Asp	Ile	Ser	Phe	His	Gly	Ile	Leu	Ser	His	Ile
			260						265					270		
	Asn	Thr	Val	Phe	Val	Leu	Arg	Ser	Lys	Glu	Gly	Leu	Leu	Asp	Val	Glu
			275					280					285			
55	Lys	Leu	Glu	Cys	Glu	Asp	Glu	Leu	Thr	Gly	Thr	Glu	Ile	Ser	Cys	Leu
			290				295					300				
	Arg	Leu	Lys	Gly	Gln	Met	Ile	Tyr	Leu	Pro	Glu	Ala	Asp	Ser	Ile	Leu
			305				310					315			320	
60	Phe	Leu	Cys	Ser	Pro	Ser	Val	Met	Asn	Leu						

Gly Leu Tyr Leu Ser Asp Ile Pro Leu His Asp Ala Thr Arg Asp Leu  
 340 345 350  
 Val Leu Leu Gly Glu Gln Phe Arg Glu Glu Tyr Lys Leu Thr Gln Glu  
 355 360 365  
 5 Leu Glu Ile Leu Thr Asp Arg Leu Gln Leu Thr Leu Arg Ala Leu Glu  
 370 375 380  
 Asp Glu Lys Lys Lys Thr Asp Thr Leu Leu Tyr Ser Val Leu Pro Pro  
 385 390 395 400  
 10 Ser Val Ala Asn Glu Leu Arg His Lys Arg Pro Val Pro Ala Lys Arg  
 405 410 415  
 Tyr Asp Asn Val Thr Ile Leu Phe Ser Gly Ile Val Gly Phe Asn Ala  
 420 425 430  
 Phe Cys Ser Lys His Ala Ser Gly Glu Gly Ala Met Lys Ile Val Asn  
 435 440 445  
 15 Leu Leu Asn Asp Leu Tyr Thr Arg Phe Asp Thr Leu Thr Asp Ser Arg  
 450 455 460  
 Lys Asn Pro Phe Val Tyr Lys Val Glu Thr Val Gly Asp Lys Tyr Met  
 465 470 475 480  
 20 Thr Val Ser Gly Leu Pro Glu Pro Cys Ile His His Ala Arg Ser Ile  
 485 490 495  
 Cys His Leu Ala Leu Asp Met Met Glu Ile Ala Gly Gln Val Gln Val  
 500 505 510  
 Asp Gly Glu Ser Val Gln Ile Thr Ile Gly Ile His Thr Gly Glu Val  
 515 520 525  
 25 Val Thr Gly Val Ile Gly Gln Arg Met Pro Arg Tyr Cys Leu Phe Gly  
 530 535 540  
 Asn Thr Val Asn Leu Thr Ser Arg Thr Glu Thr Thr Gly Glu Lys Gly  
 545 550 555 560  
 30 Lys Ile Asn Val Ser Glu Tyr Thr Tyr Arg Cys Leu Met Ser Pro Glu  
 565 570 575  
 Asn Ser Asp Pro Gln Phe His Leu Glu His Arg Gly Pro Val Ser Met  
 580 585 590  
 Lys Gly Lys Lys Glu Pro Met Gln Val Trp Phe Leu Ser Arg Lys Asn  
 595 600 605  
 35 Thr Gly Thr Glu Glu Thr Lys Gln Asp Asp Asp  
 610 615

(2) DATA TO SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 14 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide (amino acids 634-647 of hsgC1)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

Phe Thr Pro Arg Ser Arg Glu Glu Leu Pro Pro Asn Phe Pro  
 5 10

(2) DATA TO SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide (amino acids 593-614 of hsgC1)

5 Lys Gly Lys Lys Glu Pro Met Gln Val Trp Phe Leu Ser Arg Lys Asn  
5 10 15  
Thr Gly Thr Glu Thr  
20

15 (A) LENGTH: 28 basepairs  
(B) TYPE: nucleotide  
(C) STRANDEDNESS: single strands  
(D) TOPOLOGY: linear

(iv) ANTISENSE: no

AAAAGGATCC ATGTTCTGCA CGAAGCTC

28

35 (A) LENGTH: 18 basepairs  
(B) TYPE: nucleotide  
(C) STRANDEDNESS: single strans  
(D) TOPOLOGY: linear

(iv) ANTISENSE: no

ATTATGGAAG CAGGGAGG

18

55 (A) LENGTH: 28 basepairs  
(B) TYPE: nucleotide  
(C) STRANDEDNESS: single strans  
(D) TOPOLOGY: linear

(iii) HYPOTHETICAL: no

(iv) ANTISENSE: no

5 (xi) SEQUENCEDESCRIPTION: SEQ ID NO: 9:

AAAAGGATCC ATGTACGGAT TTGTGAAT

28

10 2) DATA TO SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 basepairs

(B) TYPE: nucleotide

15 (C) STRANDEDNESS: single strans

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: genomic DNA

20 (iii) HYPOTHETICAL: no

(iv) ANTISENSE: no

25 (xi) SEQUENCEDESCRIPTION: SEQ ID NO: 10:

ATGCCTGATT CCTGGGTACC

20